Carigueya, seu Marsupiale Americanum:

OR, THE

### ANATOMY

OF. AN

# OPOSSUM,

Dissed at

Gzelham-College.

### By EDWARD TYSON, M. D.


Fellow of the College of Physicians, and of the Royal Society, READER of Anatomy at Chirurgions-Hall, and PHYSICIAN to Bethlem Hospital.


#### LONDON:


Printed for Sam. Smith, and Benj. Walford, Printers to the Royal Society, at the Princes Arms in St. Paul's Church-Yard. MDCXCVIII.













w (a), and to introving Got Place (b) celetil

Carigueya, seu Marsupiale Americanum:

OR, THE

## ANATOMY

OFAN

# OPOSSUM.

And preferred to the Royal Society, by Will. Bird, Eq; and kept alive in their Repository for tome time. But of late languishing and falling from its Meat, it died. The Cause of its Death we shall see, when we come to the Dissection: But first of all, we shall take notice of the several Names given it by different Authors: Then shall give a more exact Description and Figure of its outward Shape: And last of all, we shall observe the Structure and Formation of the Parts within; especially where different from other Animals.

might

And as to the Names, I find that Georg. Margravius (a), and fo likewise Gal. Pifo (b) telle us, That 'tis called in Brofile, and on the Sca Coasts Carigueya; by fome, and in the In-land Contries, Imparitma. In Pertguay, Tai-ibi. Franc. Hernandez (c) faith, 'Tis called, Tlaquatzin, by the Indians. Antonius Herera calls it Taquatrin and the Modern Spaniards, by a Corruption, Tlaquacum; as Job. Eufeb. Nierembergius (d') informs us. Job. Per. Maffeius (e), and Caspar Barlaus (f) calls it, Cerigo. Job, Lerius (2) calls it Sarigoy or Carigoy. Joh. Stadenius (b) or Stadius, as he is Printed in Theodore de Bry, Writes it, Servoy or Serwoy. By Cardan (i), and Oviedo (k) 'tis called Chineca & alibi Chucia. In the Molucca Islands 'tis called, Zurca. In Virginia, and generally by the English, 'tis called Opossum, as by Ralp Hamor (1), and others. Job. de Laet (m), and Captain John Smith (n) writes it, Opaf-Sum. Mr. Rey (a) calls it, the Possums, as do also our common Seamen. I do not find my Litin Name given to it by any Body, but by Gefuer (p), who, that it might not be Anonymous, forms that of Semi Vulpa, quasi Simia Vulpina; or, as Aldrovandus (q) calls it, Vulpt-Simila, as if this Animal was of a middle Nature, between a fox and Ape : And, answerably, in Greek it

N. B. In my Quotations of any Anthor, if what occurs be in the fame Book and Place, I always refer to the first Letter its quoted by if otherwise, its Assinguished by different Letters, which I choose, to avoid needless Muttiplication of References.

m 14 References.

(a) Hift. Nat. Brafil. 1, 6, cap. 2, p. 222. (b) Hift. Nat. & Med. L. 5.

(c. 22, p. 323. (c) Hift. Mexican. 1, p. 1, 18, p. 330. (d) Hift. Nat. Hib. 9.

(a. 2, p. 356. (c) Hift. Indic. L. 2, p. m. aps. (f) Res Geitz ful. Mauritio in Brafilia p. m. 222. (g) Hift. Navigat. in Brafilian, c. 9, p. m. 180.

(b) Hift. Brafilian. Cap. 32, p. m. 129. (f) De Subcilitat. L. x. Oper. Thm. 2, p. 321. (l) In Summario India Occident. (l) Defeript. Virginia (m) Deferiptio India Occident. L. 2, c. 170 p. 82. (n) Vinger and Diferences in Virginia, p. 27. (c) Synopsis Animal. p. 182. (p) Hift. Anim. de Quadruped. p. 870. (q) De Quadruped. Digit. Viviparis, L. 2, c. 4, p. 223. might

might be called, Hologo Narray, or 'Angreso ribors'; though he tells us, That the Alopecopithecus and Chinesa, by Cardan and Scaliger, are taken to be different Animals.

Upon the whole, fince it is an Animal fui Generis, and in several Parts having a great Resemblance to those of different Species; I think, a Denomination might be best given to it, from that Particular, wherein its most distinguished from all others; which is that remarkable Pouch or Marsapium it has in the Besty; into which, upon any Occasion of Danger, it can receive its Young. Not that it is a Vierus, or the Young Ones are bred there (this Mistake we shall clear up, when we come to the Diffection): But Nature has only formed this Part for that Particular Vie of receiving or emitting, at Pleasure, the Young Brood, till such time as they are able to provide better for themselves.

This Consideration (it being so distinguishing a Character of this Animal from all others, that as yet we know of,) makes me most inclinable to find out some Name, that might be most expressive thereof; nor can I think of, at present, a better, than to call it, Marsupiale Americanum. If upon Observation (as very likely) there should be found different Species of these Animalia Crumenata, as Jul. Cos. Scaliger (r) calls them, by an Epi-

thee, they may be subdivided.

Bishop Wilkins (s), and Mr. Ray (s) too, in his Synaphis Animalium, places this Animal amongst the Dog Kind. Guil. Piso (b) saith tis, "Bestiola majorum Glirium Man"tanarum numero ascribenda: And Cardan saith (i),
"Mistis & India Occidentalis Chinreham & Mustelino
"Genere. I am apt to think it may be reduced to she

A 2

<sup>(</sup>r) De Subtilitate ad Cardan. Exerc. 206. p. m. 277. (s) Real Charaller, Part 2. c. s. p. 159.

Vermine Kind; but that may be better determinated, had one an Opportunity of observing the Male; for our Subject was a Female, which I am now to give a De-

fcription of.

And for the better doing this, I caused a Figure + + Fig. 1. of it to be made, exact from the Life: for those Figures that hitherto have been given of it, are very faulty; as will appear by comparing this with that of the Caragueya of Margrave (a), which is copied by Pife (b) and Johnfon (t). The Figure of Job. Eufeb. Nierembergius (d) indeed, feems to be taken from the Life, and more agreeable; but not in all Particulars exact. That in Gefner (p) is altogether Monstrous; but withal, he tells us whence he had it; " Imaginem addidi ; qualis in Tabulis Geographicis depingi folet; and mistruits it himself, where he faith, & Icon fi quid ei credendum. However, Johnston (who never fails to copy an Error, and never mends one) adds this Figure too, vid. ejuld. Tab. 58. though he owns that he had feen that of Nierembergius. The Figure of The Aldrovandus (q) is much the same with that of Gelner's.

Should one here indulge the Imagination so far, as in the Description of this Animal, to borrow its several Parts from those of different Species; one should rather seem to form a Chimerical Monster, than to describe a real Animal. Yet we find the best Zoographers thus to

please themselves in their Accounts of it.

Animal est Tlaquatzin Canis parvi formà & magnitudine, saith Hermandez (c). Animal est Catuli vel Mediocris felu magnitudine, as Margrave (a). Pars anterior Vulpem, posterior verò Simiam representabat, nist quòd Pedes essingit Humanos, Aures antem babet Noctue, saith

<sup>(</sup>t) Hist. Animal Quadruped. Tab. 63.

Vincentius Pinzonus (u). Monstrosum illud Animat Vulpino Rostro, Cercapitheci Cauda, Vespertilionis Auribus, Manibus Humanis, Pedibus Simiam amulans, faith Peter Martyr (w). Corpore verd ac Pedibus Melis, the Mexican History (c). Videtur autem boc Animal intermedium effe inter Leporem ac Vulpem, sed alterius generis ab utroque ob Crumenan, faith Cardan (x). Eft Viverre facie ac magnitudine, Capite Vulpino, faith Jul. Caf. Scaliger (r). Opassim Capite Porcellum, Cauda Glirem referens, Mole modicum Felem, faith Job. de Laer (m). Barbam Felinam habet, faith Pife (b). Cauda Colubrina fimilis, Hernandez (c). I shall omit more Instances which might be given. Nor is it that I do disapprove of these Allusions upon the whole; but when they call it Animal Monstrofum, as P. Martyr (w); or Prodigiosum, as Vinc. Pinzonus (u), and Hieron. Benzon. (); I think's is only our Ignorance makes the Admiration, and that Admiraration forms the Monster; for Nature, in her regular Adings, produces no fuch Species of Animals.

In our Description of this Animal, we will first give the Dimensions of this we dissected; and then proceed to other Particulars. As from the Extremity of the Nose, to the Tip of the Tail, it measured One and Thirty Inches: The Length of the Head was Six Inches: The Tail was One Foot long: The Neck and the Body was the Complement of the First Dimension: The Girth of the Body, now dead, was Fisteen Inches and an Half; when alive and well, it seemed much thicker: The Fore-Legs were Six Inches long; the Hinder-Legs but Four Inches and an Half: The Girth of the Tail, near the Root, was Three Inches; near the Tip but One Inches

<sup>(</sup>a) Navigatio inter Novi Orbis Scriptr. p. 86. (b) Ocean Decad. 1. lib. 9. (x) De Rerum Varietate, Cap. 33. p. m. 101. (7) Hieron. Benzon. Hift. Novi Orbis, Lib. 2. p. 52. apud Theodor. de Ery Americ. pars quinta.

The Head, about the Ears was largest; measuring on the Forehead, from one Ear to the other, Three Inches; thence gradually tapering towards the Nose, and more resembling that of a Fig than a Fox: The Aperture of the Eyelids were not Horizontal, but lying in a strait Line from the Ears to the Nose, and not large: The Ears were about One loch and an Half long; not sharp, but of a roundish Figure: The Risks of his Month, from the Corner on One Side, to the End of the Nose, meafured Two Inches and an Half.

Thefe Merfures, in feveral Particulars, I find different from those in other Authors: But I cannot but think they have made several Mistakes, for want of a more exact Enquiry and Observation. Thus Margrave (a), and, from him, others makes the Fore-Legs thorser than the Hinder; " Crura due anteriora (faith he) breviora, mimirium quodlibet tres digitos longum; Pofteriora paulo plus quatuer : In our Subject I find the contrary; tho, as we shall observe in the Sceleton, the Bones of the Fore-Legs are shorter than whole of the Hinder. He adds, " Pedes Anteriores quinque digitis Instar manus constant, " unquibus albis instar Avium, Curvis: Pasteriores longio-" res, uti in Cercopithecis effe folent, item quinque digitis at manus. But here we find the Fore-Feet to have Five long Claws or Fingers, equally ranging with one another; and a hooked Wail at the End of each Finger. But the Hinder-Legs are far differently formed; for here we obferve but Four Fingers armed with hooked Nails, and a perfect Thumb, fet off at a Distance from the Range of the other Fingers; and as in a Humane Body, this Thumb was fhorter than the other Fingers; and had not a hooked or curved prominent Nail, but a tender flat one, as in our || Figure is represented.

|| Tab. 1.

This Contrivance of the Feet and the Nails, and the Fore-Feet being longer than the Hinder, and the Hinder being formed with a Thumb like a humane Hand : feems very advantageous to this Animal as to its way of living, and getting its Food : For what it feems to be most fond of, is, Poultry and Birds; not but that it cuts other things too. But for the Preying upon thefe, 'tis very nimble in climbing up Trees; "Scandit Arbores incredi"bili pernicitate, faith the Mexican History (c); and the same is confirmed by all. Now having the Hisder Feet formed like an Hand, and the Four Fingers Armed with hooked Nails, it may takethe bester Hold in railing its Body upa Tree; and the Fore-Legs being longer, will make the larger Stretch in Climbing up; the Mails. being hooked and flrong, will take the greater Hold. Befides, as we shall observe, having the Advantage of twifting its Tail about any Stick or Bough it lights upon. and being able, by that Means, to fulpend the whole-Weight of its Body; when it has a mind to raife it felf to the Bough its Tail is twifted about, thefe Hinder-Legs, being formed like Hands, will the better take hold of it,. than if they had no Thumbs.

These Fingers, Toes, or Claws were maked, without Hair; the Skin looking of a Reddish Colour here. They were about an Inch long, and the Thumbs almost as long, but set lower, as I faid. The Palmes of all, especially if dilated, as it does in Climbing, were large; but to contrived, as to be able to be contracted, as in walking; but that they might here be better fecured from Injury, I find at the fetting on of each Toe, in the Palmes, a. Protuberant, Fleshy, and almost Cartilaginous Body.

In feeding its felf, it makes use of the Fore Feet in bringing the Food to his Mouth, as do the Monkey and

Squirril Kind.

As the Toes, so likewise the Tail was without Hair, only for a little way near the setting on; and was most remarkable; being tapering from the Root towards the Tip, and covered with a regular Order of small Whitish Scales; which, for the most part, were all oblongish Hexigons, Persimilis Colubrina, saith the Mexican History (c): But with this difference, that in a Snake's Skin part of one Scale juts over another; but here each Scale appears wholly in view, and between each one may observe a little of the Skin or Membrane in which they are fixed. The Colour of these Scales, makes the Tail to appear Whitish, though the Skin seems of a Darker Colour.

As the Feet and Tail, so likewise the Ears were bare, and without Hair: Auriculis tenuissimis, mollissimifq; ac pene translucentibus, saith the Mexican History (c). And although thus foft and flender; and in Colour and Substance almost resembling the Membrane of a Bat's Wing, yet they were creek, not acuminated or running to a Point, as described by some; but, as in our Icon is represented, more of a Circular or Oval Figure: They were so slender and soft, that here I could not perceive that Cartilaginous Body, which usually is to be met with in the Structure of this Part, in most other Animals; but as if it was void almost of it, to be formed only by a Duplication of this tender Membrane or Skin; or if it did enjoy a Cartilage, as likely, 'twas much finer than in most other Animals. The Concha or Passage to the Porus Auditorius was very Capacious: But 'twas observed, that when our Subject began to grow ill, the Verge or Rime of the outward Ear feem'd to be crimp'd; and when it died, to be so shrivel'd, as if burnt up, not making a smooth, but jagged Edge.

The general Vogue of almost all Naturalists, resembling the Face and Head of this Animal, to that of a Fox,

would

would much incline me, could I find but a tollerable Agreement, to acquiesce herein: But by comparing both together, I see but very little Likeness. A Fex. tis true, has a large Forehead, and a flender Roftrum : but in a Fox it does not gradually grow Taper (as in our Subject, from the Ears to the Note); But here the Forehead is Prominent, and somewhat Oval or Circular: then makes a remarkable Break; whence is protruded a flender Rostrum. I much more think, in this Particular, our Animal refembles that of a Hog: But Comparisons being so delusive, so un-satisfactory to an inquisitive Mind; and in all Natural History not meeting with more, and to so little purpose, as in the Description of this Animal; I should rather think it far more adviscable, to avoid the seducing the Mind into an Error hereby; than at the same time, when 'tis pretended to inform, to mil-guide the Imagination by a mistaken and ill-chosen Simile.

I shall only add, that the Upper-Jaw was somewhat longer than the Under: The Nostrils were large; Nares babet patentes, saith Margrave (a): The Eyes Black, small, vivid and exerted, when alive; now dead, very much sungk: The Neck was short: The Breast was broad; and the Shape of the whole is best appre-

hended by confulting our Figure thereof.

It had Mustacio's like a Cat; Barbam babet Felinam, faith Margrave (a); but his Pidure of it is very much mistaken. The Fur upon the Face is shorter and whiter than the rest of the Body: The Mexico History (c) describes it Rostro tenui, prolixo & depili; but this must be only meant comparatively. On the Back and Sides the Fur was of an Ash Colour, or dappel'd with Black Hair in Spots, intermixt with White, especially on the Back; on the Belly 'twas more of an Umber Colour, and of a darker on the Legs. The longest Hairs, which were B

stronger and courser than the rest, measured Three Inches; being White towards the Ends: though the Mexican History saith the contrary. Pilo longo & candido, sed circa extrema susce & nigro, saith Hermandez (c). Job. Euseb. Nierembergius (d), in his Figure (which is much to be preferred before the others) represents him very Shocky, and, as it were, with Curled Hair. If what Margrave tells us be true, his Subject was different from ours; Pili autem Capitis (saith he (a) Colli inferius cum fine Ventris & Cauda inferius prope exortum, sant slavi secundum longitudinem autem Capitis, per Oculos & medium Capitis tendit ampla nigra stria. But the Colour of the Fur may vary in different Subjects.

But we will have now done with the Outward Parts, to proceed to the Inward: But that we must first of all take Notice of that most remarkable Part, that no other Species of Animals enjoys, as I know of, but this; that is neither Inward, nor Outward, but a Medium between both; I mean that admirable contrived Pouch or Marsu.

pium that it has in the hinder Belly.

For at the Bottom of the Belly, in the middle, between the Two Hinder-Legs, we observed a Slit || or || Tab. 1.

Aperture, moderately extended about Two Inches long; Fig. 2. and capable of a larger Extension by dilating it with ones Fingers, even when it is alive. John Stadius (h) saith, in alvo fissuram habet ad dimidiæ Spithamæ longitudinem: Or, as 'tis rendred by Gesner (p) ad sex digitos sere scissum: Possibly 'tis a little over stretcht; however, it can so exactly close and contract it, that the Eye does not readily discover it, till dilated by the Fingers. "Bur-" sæ os ita clauditur, ut non appareat, nist duobus digitis ab invicion distendatur, saith Margrave (a). Tanta æ-" qualitate & vi, ut coaluisse cutim omnino putes, saith the Mexican History (c).

This is so surprising a Structure, that all Zoographers do mention it with the greatest Admiration; and yet their Curiosity has not been so great, but several Mistakes they have been guilty of, in the Accounts they

have given about it,

Mirum autem Animal (saith Margrave (a) nam in infimo Ventre prope Crura posteriora Pellis ejus dupla est, &
exterior rima duos & semis digitos longa, facita; quasi
Bursam, quem Brasiliani vocant, Tambeio, Pomi Aurantii
majoris capacem. Est autem Bursa bæc intus Pilosa. All
of this, I think, is very just and true: For at the Place
he mentions, we observed an Aperture, or Slit, much of
the Largeness he describes, where there is of each Side,
a Reduplication of the Skin inwards, which forms a Bag
of the Capacity he mentions, and of a greater upon Occasion, and Hairy too. But these Hairs here are so thinly set, that by no means they cover it, but almost every where you may observe the Skin.

Gul. Pifo (b) has much the same Description, but calls this Bursa, Mantica. Cardan (x) calls it Crumena. Oviedo (k) calls it Ventrale Marsupium. By Joh. de Laet 'tis stiled, Ascus sub Ventre pendulus (z). By Fr. Ximenez, or his Translator (aa) Pelliceus Saccus. By Jul. Cass. Scaliger (r), Scortum subventrale. By Peter Martyr (w), Uterus Exterior. By Hieron. Benzon. (y), Venter alter. By Captain John Smith (n) 'tis called a Bag. The Name we shall use for it, will be, the Marsupium or

Pouch.

Herein all agree, that the use of this Bag, Pouch, or Marsupium, is for the Preservation of the Young Ones, and securing them upon any Occasion of Danger: Qua-

E (x) Joh. de Laet. descrips. Indix Occident. lib. 15. cap. 5. p. 551(aa) Vide Joh. de Laet. ibid. lib. 5. cap. 4. p. 232.

ternos, quinofve parit Catulos (faith Hernandez (c) quos Utero conceptos, editofq; in lucem, Alvi Capacitate quadam, dum adhuc parvuli sunt, claudit ac servat. Peter Martyr (w) tells us, Quod natos fibi Catulos circumfert, quocunque profiscatur, Utero exteriore in modum magnæ Crumene dependente, &c. quo à venatoribus vel alias à cæteris violentis ac rapacibus Animalibus natos liberet. illos fecum asportando: nunquam autem illos emittere dicitur, nifi aut recreandi, vel lactandi Gratia, donec fibi vi-Elum per se queritare dedicerint. Oviedus (k) adds, Quod Bestia bæc ritu mustelarum noctu domos ingrediatur, necanfq; Gallinas, earum sanguinem tantum modo sugat : imo illic Ventrale Marsupium aperit : filiosq; dimittit, ut & ipfi ad Gallinaceum sanguinem sorbendum assuescant : interim fi aliquem strepatum senserit, illico natos in crumenam recipiens, fugam arripit. And Job. Stadius (b) owns he has taken out the Young Ones hence; Aliquoties ealdem ipse venatus sum, & Catulos ex finu ipsemet deprompsi. This Particular, I think, is evident by the Contession of all; and thererefore shall not use more Quotations to confirm it: And Nierembergius (d), on this Account, in his Picture of this Animal, represents the Young Ones as crawling out. But when they tell us, there is no other Vterw : But that the Fætus are formed here. and nourished here; and this too, when they own they have diffeded this Animal; this is too great a Mistake to be longer Propagated, and from Autophe we shall demonstrate the contrary, when we come to the Diffection.

But fince the great and wife Defign of Nature, in contriving this Part, is for the Prefervation of the Young Ones, in receiving and emitting them as there is Occasion; we will contemplate and admire the admirable Structure and Artifice she uses in forming and adopting it so suitably for this End.

We will therefore here remark the Structure of the Muscles, which, like Strings, do serve to open and shut this Pouch. But in the doing this, we must first describe those Two remarkable Bones this Animal has, more than is to be met with in any Sceleton besides, which are of great Use and Service herein: And, from their Office, I shall take leave to call them, Ossa Marsupialia, or Janitores Marsupii. I do not find they have been yet taken notice of by any; and though they do belong to the Sceleton, yet I can't avoid their Description here, because of the Muscles that are inserted to them, that do serve to open the Pouch.

|| Tab. 1.

+ 6.

\* d. e.

Hd.

te.

These Offa Marsupialia | or Janitores Marsupii (as I call them) are Two strong Bones, so fastened to the upper and inward Edge of the Offa Pubis, that at their Basis here, they touch'd one another, just at the Coalition t of the Bones that forms the Offa Pubis . The other Extreme of these Bones were so distant from one another, that it measured Two Inches and an Half. The Basis tof these Bones, where join'd to the Offa Pubis, was half an Inch broad, having Two Heads; the larger ! lying near the Coalition of the Offa Pubu, and the leffer † towards the Os Coxendicis; having in the middle a Sinus, into which was received a Protuberance of the Offa Pubu ? By which Contrivance it appears, there can be no Motion of these Bones, nearer or farther from one another, but that they must stand always at an equal Distance. Nor did I, upon trial, find it otherwise; but observed, they were capable of a small Motion inwards towards the Spine, and outwards from it. These Bones, as they afcended from the Os Pubis, grew slenderer, being about the Middle but a quarter of an Inch broad; and they were each, Two Inches long.

To each of these Bones there were bestowed Four Pair of Muscles: There was another Pair that did run over

them

them; to which these Bones did perform the Office of a

The First Pair of Muscles (i.e. which first came to be diffected, upon the Pronation of the Animal, and from its Figure, I shall call, Triangularis) arises Fleshy from the whole Length of the internal Side of these Bones, and inferted their opposite Tendens on each side of the Rima. or Aperture of the Marsupium. Under part of these Muscles, lay another, or a Second Pair; flat and thin; having their Origin from the upper Part of the internal Side of the Offa Marsupialia and inserting their opposite Tendons a little above the Tendons of the former Muscles. the Tendence or Direction of the Muscular Fibres of this Pair, in Relation to the First, made a Decustation. The Third Pair of Muscles we shall take notice of, had their Rife from the Fore-part of the Bafis of these Bones, where they were jointed to the Os Pubis; and were afterwards inferted into the Linea Aspera of the Thigh-Bone. The Fourth Pair did arise from the external Side of these Bones near the Bafis, and are inferted into the Fore Part of the Thigh-Bone near the middle.

The Last Pair of Muscles I hinted at, (to which I thought these Bones might perform the Office of a Troreblea, or Pully) arises more immediately from the Marsupium or Ponch it self: For spreading their Muscular Fibres all over this Bag, as they issue from it, by joining their Fibres together; they more remarkably form a folid Muscle; which of each side passing over the middle of these Bones, (i. e. in the prone Posture we are dissecting it) at length were inserted into the Spine of the Os

Ilii.

By considering the Structure of these Muscles, and what must be the Effect of their Adion or Contraction; one cannot but think the Two First must serve towards the Dilatation or opening this Marsupium or Pouch: For these

these Bones are a Fulciment or Basis; their Articulation will not admit of a Contraction inwards or nearer to one another; wherefore, when the First and Second Pair of Muscles act or contract, they must necessarily open or dilate the Mouth of the Marsupium or Pouch. The Third and Fourth Pair of these Muscles may serve to extend these Bones outwards; so that when this Animal baurs by its Tail (as it does frequently) the Weight of the Fatus in this Pouch by this means will not press so much upon the inward Viscera. The Fifth and Last Pair, as they may ferve to dilate the Capacity of the Pouch it felf, fo. likewife may ferve the better to suspend its Weight, when the Animal is prono Capite, and if it gravitates too much. they may retract it up, and the easier, because pasting over thefe Bones like a Pully, their force is more augmented.

The Antagonist to these Muscles is, the Sphinder Marfupis; an oval Series of strong, sleshy Fibres, which serve to constringe and close the Orifice of the Pouch; which it does so perfectly (as I have already observed), that one would think the Skin here not to be slit; nor can the Orifice be observed till you have dilated it with your Fingers.

Nature's Contrivance therefore in placing this Ponebhere, in this Hinder Part of the Body, is very great; her Mechanisme in forming these Two Bones, the Janitores Marsupii, which no Sceleton besides has; and so artfully furnishing them with these Muscles, is most admirable; that with the Philosopher, there is none but must own. Dec yeaperpel.

The Pouch or Marsupium it self, was a Membranous Body, not very thick, tho' consisting of several Coats, and is to be reduced into the Class of the Vesiculous Parts of the Body; which according to my Notion, are part Muscles, part Glands, and do perform the Office of both,

Motion ..

Motion and Secretion : for the Concave or Hollow + of + Tab. 1. this Pouch (as I have remark'd) was somewhat Hairy; Fig. 3. and at feveral Places I could observe them matted or cling'd together by a Tellowift Substance, which did onze out of the Cutaneous Glands there; as under the Armpits in a Man, it is observed. This Liquor thus emptied into the Pouch from the Glandulous Coat, I found was strong Scented, and had more of the Peculiar Fator of this Animal, than any part belides; being no ways grateful, but unpleasant to the Smell, as has been observed of this Creature, when alive : Fætet Animal instar Vulpis vel Martis, faith Margrave (a). Vulpeculis Hi-Spaniensibus sunt similes sed minores, & longe graveolentiores, faith 7ob. de Laet (2). But after the Skin with the Pouch had been kept for some Days, and was grown dry, I found fo great an Alteration here in the Smell, that what before was fo difagreeable, now was become a perfect Perfume, and finelt altogether like Musk; which made me call to mind what formerly I had remarked (bb) of these Scent-Bags in other Animals; that in a Weafel, which Stinks like a Poleicat, by spreading this Fæted Liquor contained in the Scent Bogs (as I call them) on a Paper, and fo letting it dry, it became a grateful Perfume, &c. wherefore afterwards, in my De-Scription and Anatomy of the Tajacu(cc) (a Mexican Wild-Hog); when I came to diffect it, fome Days after it had been dead; and the Virus or horred Fator of the Liquor contained in the Scent-Bag (which was a large Gland on the Back) had fpent its felf, and it became an agreeable Perfume: Upon this Consideration, I took the Liberty (it may be, a too great an one) to call it Moschiferus; Bady, nor very thick, tho confi

though

<sup>(</sup>bb) Philos. Transact. No. 144. p. 39. (cc) Philosoph. Transact. No. 153. p. 379.

though the general Consent of all Authors had branded it with the Note of a Fatid stinking Animal. But at the same time, I instanced, that in the richest Persumes we have, as Musk, Civet, and Ambergriese, the same is observed; and that Passage I quoted in Gul. Piss (dd) con-

cerning Ambergriese, is very remarkable.

Upon looking over this part of my Account of the Tab jacu, I find this Passage (cc): " And I am apt to think, "'twas by removing thefe Scent-Bags, rather than ta-"king away the Kidneys, that they made the Sarager "edible; which otherwise stank to much, that the " Barbarous Nations refused them; as out of Lerius. " Job. Faber (ee) takes notice. Job. Lerius his Words are these, as he is printed in Theodore de Bry (ff) alia etiam invenitur (Bestia) quam Saragoy appellant, que ob fetorem a Barbaris non comeditur, nos autem quibusdam excoriatis, & detracta Renum pinguedine, unde fator ille manebat, fine fastidio edimus, carne enim cum tenera, tum optima eft. And I find my Conjecture was not amis: for although then I knew not where this Scent-Bag was placed, yet now I find it was the Marsupium, or Ponch : and that Lerius was altogether mistaken in attributing this Fator to the Kidneys, or the Fat about them o for I could not smell any thing ungrateful, or ill-scented there.

Upon this Occasion, I can't but make a farther Remark upon the Scent-Bag of the Musk-Deer; which is likewise placed in the Belly near the Navil, and makes a large Cod. Lucas Schröckius (gg) has wrote a distinct. Treatise about it; to whom I shall refer my Reader. But though there is this Analogy between our Possum and

C

<sup>(</sup>dd) De Indiz utriusq; re Nat. & Med. lib. 1. p. m. 17. (ee) Histori Mexican. p. 658. (ff) Americ. part tertia p. 180. (st) Historia Moschi-

the Musk Deer, that both have their Scent-Bags placed in the Belly, and they have an Aperture there; yet the grand Use of the Pouch of the Possim is very different,

as we have already feen.

Having therefore mentioned the Glandulous Coat of the Marsupium (on whose Account it may be reckoned as a Scent-Bag) I must here take notice that it had likewise a Muscular Coat, besides those several other Muscles bestowed upon it, which we have observed already, that gave it Motion. It had likewise a Vascular Coat too, being plentifully irrigated with Blood-Vessels, especially by Two large Branches that came from the upper part of the Thorax, and might be reckoned the Mammaria, as they are stilled in other Animals. This Pouch was saftened by several Membranes to the Muscles of the Abdomen and the Skin; but so as I could separate it for the most part, with my Fingers.

But here I find, in this Marsupium or Pouch, they place the Mamma or Teats; and they tell very odd Stories about it: I will only relate what they say of it, and what I at present observed, or rather, did not observe.

Job. Petrus Maffeisu (e) makes this Peuch, not single, but double, he places the Mamma here, and the Young Ones to be so fixt to them, as if they almost grew there. Illud autem mirum in Cerigonibus, ex ejus alvo dua dependent veluti Mantica, in iis catulos circumsers. O quidem adea pertinaciter sun quemq; Uberi assisso, ut a perpetuo sulla non avellantur, antequam ad pastum ipsi per se progradi valeant. And much the same has Casp. Barlaus (f) Cerigones (saith ho) Kulpis magnitudine, insolito spectaculo alvum ostentant, è qua dua veluti Mantica dependent, quibus catulos gestant, tam valido suctu Uberibus adbarentes, ut non antea demittunt, quam adultiores ad pabulum ipsi excurrere valgant. Gul. Piso (b) out does them both; sor he makes them not only Nourished, but Formed here:

here: Ex reiteratis (laith he) borum Animalium fectionibus alium non invenimus Vterum, præter bane Burfam, in qua femen Concipitur & Catuli formantur. Quos deinde Quinos vel fenos fimul circumfert, mobiles, perfectos, fed depiles adeog: pertinaciter Oberibus affixos, ut a perpetuo fullu vix avellantur, antequam permittente Matre ad paftum ipft egrediuntur; unde redientes denno, uterum maternum pro lubitu ingrediuntur. So Job. Stadius, as I find him Printed in Theodor de Bry (bb) In Alvo (faith he) fissuram babet ad dimidiæ spithamæ longitudinem,intra eam alia Cutis subest, nec enim alvus biat aut fatbifeit. in eo finu sunt Obera, quocunque obambulat proles secum gestat. So likewise Peter Bembus (ii), writing of the new discovered Islands, faith, Animal ea sylva untriunt Cuniculi magnitudine, Gallinis infestissimum, enjus quidem fæmina loculum babet e Pelle Utero amexum, quafe Uterum alterum fæcundum Vberibus, in quo Catulos fecum geftat. emittitg; cum vult. And our Country-man, Captain Smith (n) feems of this Opinion too; where he faith. Under ber Belly she bath a Bag, wherein she lodges, carrieth and suckleth her Toung. But Gillius, as he is quoted by Gefner (p), tells us, That non ex receptacule prodeunt. nift cum lac sugunt. So likewise Vincent. Pinzonius (u) Nec unquam exeunt Crumenam, nifi quum sugunt ; which intimates, That the Teats are not placed here, fince they must go out to fuck and and promise in visite & and

I must consets, upon what Observation I could make, I did not find any Teats here; nor indeed did I find them in the outward Skin; as is usual in other Multiparous Animals. Possibly this Subject never had a Litter; so for want of drawing, they might be less; so as to escape

<sup>(</sup>bb) Americ. Part Tertia. Joh. Stadii Hist. Brasilian. cap. 32. p. 129.

our View, for the present : But in another Subject, I

doubt not but that they may be discovered.

But this Bag or Pouch is not only appropriated to the Female, as one would guess by what P. Bembus (ii) faid before: But if we may believe G. Pifo (b), the Male has one too : " Mar fæmellæ plane fimilio, &c. (faith he) & quod notatu dignum, manticam (licet a famella diversam) babet ; qua alternatis vicibus Catulos quoque circumfert ; ficut avis mas amore pullorum, fæmellam ab incubatu subinde liberans. I could have wished he had given us an Account of the diversity of the Pouch in the Male, from that of the Female; and had one an Opportunity of Diffecting a Male, it would much illustrate the Hiftory of this Animal. So likewise the Author of the Present State of his Majesty's Isles and Territories in America, (pag. 138.) tells us, That the Male bas such another Purfe under bis Belly, and takes his turn to carry the Toung Ones, to ease the Female. But more of the Male, when I shall come to mention the Tai-ibi.

This Contrivance of Nature for securing the Toung Ones from any Danger, till they are able to shift for themselves, I think, is not to be parallel'd in any Species of Animals, at least of the Quadruped Kind, besides. Not that the is wanting in abundantly providing for their Preservation; but the pleases her self in using infinite Variety in attaining the same End. Nor is there wanting Instances enough to evince it: What most reaches, and comes up to our Subject is, what I find recorded in Oppianus (kk) in his excellent Poem of Fishes: For in his Halieuticks, describing the Philostorgia of Fishes, having mentioned the Dolphin, he comes to the Dog-fish, and tells us, that upon any Storm or Danger, if

pursued, the Toung Ones run into the Mother's Belly; and when the Fright and Danger is over, they come out again. I shall not think much to transcribe his Verses; which were so admired by Antoninus, the present Emperor of Rome, to whom he made the Dedication, that not only for them, he revoked the Banishment of his Father; but presented bim likewise with a Golden Statera for each Verse; whence they are called Golden Verses: Which, according to Suidas's Computation, came to Twenty Thousand. Suidas (II) his Words are these: 'Anaproching dural προς ένα είχου ἐν μέθρω, εαθηρα χρυσών, ήγων νόμισμα. Το καθείν αυθόν δτὶ πῶν, νομίσμαθ. μυριάδας 6.

And upon this Occasion, Oppian not only instances in the Dog fish, but in the Squatina, and Glaucus too: But there are different Receptacles he does assign, that these Two latter do make use of, for the receiving their Toung, in Case of Danger. Oppian's (kk) Verses are

thefe:

Τρούσι δ' αῦ δελφίς μὲν ἀρος δύοι φιλότηπη Παύδων ῶς ἢ ὰ ἀλλοι ἐὰν γίν Β. ἀμφιέπωσι. Θαῦμα δ' ἀλιπλάγκτοιο κυνός τόδε. τῆ γδ ἐπου) Τέκνα νεοδλας ῆ, καὶ σφιν σάκ Β. ἔπλετο μήτης. Αλλ' ὅτι τας δήσωπ, τάτ ἀμπιτα δείματ ἔμπις. Εν πόντω, τότι παύδας ἐνὶ λαγόνεων εθέκτο Αὐτίω εἰσθμίω, ἀμτίω όδου, ἐνθεν όλιαδον Γοινόμενοι. τοῦσι δὲ πάνον μογίωσα πιρ ἔμπις. Αμπιτάνος τέτλημε, πάλιν δ' ἀπεχάνατο παύδας Σπλάγχνοις ὰλ δ' ἀνέπκεν, ὅτ ἀμπιτάνου πρόδοιο. Τοῦω ὰ ρίνη τεκέων πορσύνε) ἀλκίω, Αλλ' ἐκ ἐς νηδοιὸ κοίνη δύσις, οἶα κύνεωπν.

<sup>(11)</sup> In Voce Oppianus.

Αλλα οἱ ἐν πλουρίπου ἐποσφαρις αμφοτήρωθεν
Εἰπν ἐπο περύρων, οἶη γένις ἰχθύσιν άλλοις.
Τῆπο ἀτυζομένων, τεκέων φόδον ἀμφικαλθπέω.
"Αλλοι δ' αῶθ' ἐὰ τέκνα διὰ τόμια παρβήσοντα Δεξάμενοι ρύον), ἀτ' ἐς δόμιον ἡὲ καλιήν.
Οἴον ἀλ ὰ γλαῶκῶι, ὅς ἔξοχα τέκν ἀραπάζω Παντων, δοποι ἐαπν ἐν ἰχθύσιν ώστουῶες.
Κῶνῶι ρῶ μέμνω τι παρήμενῶι, ὅρρα γένων)
Παῦλες ἐποκάδιοι, καὶ σριν παρανέκα) ἀιελ.
Τὸς δ'ὅτι καν τρομέρντας ίδη κρατιρώτες ον ἰκθιώ,
Αμρικανών κατάδικτο καὶ ὰ τόμια, μέσρα κα δώμα Χάοπ) τόπ δ' αὕτις ἀνέπους λλικανόνον.

Which an ingenious Friend of mine has thus translated: And indeed, the whole Poem is so fine, so noble and rich a Thought; that if the Translation of it was attempted by some good Genius, he would find Charms enough in it to raise his Fancy; and a Subject Worthy his greatest Skill.

None to their Young so kind as Dolphins are;
Tho' other Fish of theirs, express their Care.
This in the Dog-fish, we with Wonder see;
To whom her trembling Whelps from Danger slee.
For when a dreadful Storm imbroils the Deep,
Within her Bowels, they for Refuge creep;
They pass directly to her Womb; the Way
By which excluded, first they saw the Day.
She, tho' distended, and in torturing Pains,
The bulky Burden patiently sustains,
While the loud Terrors of the Tempest last:
But when the Danger, and their Fears are past,
Strait from her suffering Womb, the Whelps retreat,
And she her Labour, they their Birth repeat.

The Squatina do's the same Love betray,
As Dog-fish do, but in a different way.

Nature has Bags on either side prepar'd

Beneath her Gills; where she do's hide and guard

Her frighted Young; with sudden Danger scar'd.

Some thro' the Parent's gaping Throat descend; Whom as a House, or Nest, their Maws desend. The Glaucus, for Assection, is by none Of all the Fish Oviparous out-done.
Close by her Spawn, she does unwearied stay, Her Off spring to protect from Fish of Prey. And when, with tender Fins, they strike the Tide; She with them swims, their constant Guard and Guide; For if it happens, that the trembling Fry Do some Voracious Enemy descry, Opening her Jaws, she kindly does devour Her Young; to save them from the Invaders Power-Ibeir Fears removed, she spues them up; to ease Her Labour, and restores them to the Seas.

I must consess, I would not expect in a Poet, that Nicety and Exactness of Natural History, as in a Philosopher, who is not to give a loose to his Imagination, but truly to relate Matter of Fact. But this Particular of the Dog fish, with me, bears the more Resemblance of Truth; for formerly, dissecting a Fish of this Bind; that was a Female; as it was surprising to me at the same time, to observe here the gradual Formation of several Fetus; some just beginning to be formed, others an Inch long or Two; others Four or Five; some Nine or Ten, and fit for Birth: So what comes most to our Business was, That near the Exit of the Pudendum of each side, I observed a Foramen or Hole that was capable.

the second of the second

ble of Extension, and readily enough would admit my Finger, which led into the Cavity of the Abdomen or Belly it self; and not into any Bag, or the Vierus, or any other Part. Besides, in the Abdomen I found a Quantity of Water, which I could not but think was let in this Way. So that if upon Observation at any time, there should be sound loose in the Cavity of the Belly of this Fish, a Toung one; there would be no Reason to mistrust the Relation, since here are Two Doors to let them in and out.

Ælian (mm) relates the same Story of the Dog-fish: his Words, as they are translated by P. Gillius, are thele: Si quis eorum (speaking of the Young Ones) timeat, ingreditur rursus per Genitalia in Ventrem Matris; ubi timor abierit, is prodit tanquam rursus editus. And in the preceding Chapter he tells the same (as our Poet does) of the Glaucus: And that Passage of Zeza, as 'ris quoted by the Learned Conrad. Rittershusius (nn) is very express: Γλαύκος, κύων, ή γαλεός, Βαλάπιοι ίγθύες έπεργομένε δείμα ! Θ. τοίς τίχνοις τοίς οίκείοις, δ γλαύκ Θ. μέν κ, γαλεός τε σόμαλι συγκρύπθοι, δ κόων ή κρύπθοι ταύλα πάλιν εν τη νηδύϊ, καὶ πάλιν ταῦλα γενοῦ το ρόδο παρελθόν [ ... Nay, Aristottle (00) himself acknowledges the same thing of the Galeus, which is of the Dog Kind: 'Or wer ar artor 2a-प्रका मुद्रा दिवार्वन मुत्री है और राजीवा बोद दिवारिंद राउँद महामार्थंद, में वी piva, nai vaeyas. Which Jul. Caf. Scaliger thus translates, Cæterum Mustelorum genera & emittunt & admittunt intra se Catulos; item Squatina & Torpedines. And the like he (pp) affirms of the Dolphin and Porpois : Kleis-Smorlas 3 ra recra purea cirla. But after all, if what is thus related of these Fishes should prove but a vulgar

<sup>(</sup>mm) Hift. Animal. l. 1. c. 18. (mn) Comment. in lib. 1. Halieut. p. m. 214. (o) Hift. Animal lib. 6. cap. 8. p. 677. (pp) Hift. Animal. lib. 6. cap. 10. p. m. 682.

By way one of Louis took a flash a Man-seling fift, and who Bloom onifle of Look, outstoo much grainty) a foot flower in Englice half of the flat of specific and for the seling to some on for the selection of some of selection of the selection of selec

Error, 'tis one of a very Ancient Date; and it is high time it was removed: and if there should be any Truth in these Stories, for the gaining a greater Belief thereto, 'tis requisite that 'twas supported by some more evident Proof, and confirmed by later Observation. But what we mention of our Quadruped, the Possum, is notoriously known and agreed on of all Hands; to whom, after this Digression, (too large an one it may be) we shall now return; and shall therefore be more concise in the Description of the other Parts; and shall chiefly take Notice of those only, wherein I find something remarkable, and different from the common Make of the same Parts in other Animals.

And for this Reason I have very little to say of those Parts in the Thorax, I only observed, that the Lungs had Three Lobes on one side; and but One on the other: But this One, was as large as the other Three. They were soft and spongy, and easily dilated, and large proporti-

onably to the Animal.

The Heart was included in a Pericardium, as usually; but the Heart it self I thought proportionably larger, in respect to the Bulk of the Body, than is commonly; nor was its Cone so sharp, but rather more obtuse. It had

Two Auricles, and Two Ventricles.

About the Throat there were large Glandulæ Maxillares. The Tongue was a little above Three Inches Long,
about Three Quarters of an Inch broad; 'twas rough,
having several Protuberances, whose Points looked inwards. Lingua longa (saith) Margravius (a) quam tamen os aperiens & morsum minitans non exserit, quamvis
possit, sed versus posteriora attollit. The Voice or Noise
it made, was a little Growling.

But meeting, or at least not observing any thing farther extraordinary here, we shall hasten to the Abdomen,

D

where

where we shall find more Matter to excite our Admira-

tion, and please our Curiosity.

The Abdomen or Belly was divided from the Thorax or Breast, by a large, strong, sleshy Diaphragm, for (as we shall observe in the Sceleton) the Thorax near the Throat was small; then gradually, as it descendeth, it enlarges its Capacity; fo that here, where the Diaphragm was fastened, its Compass was very great and large; which might be rendred fo the more, by reason it often hangs by its Tail; and when it does fo, the Viscera in the Abdomen can't but press upon it. But that they might not too much, to the Injury of the Animal; we shall see what Provision Nature has made for it, by her

great Contrivance in suspending the Intestines.

But we must observe and describe the + Stomach, or + Tab. 2. Ventricle, which was feated under the Diaphragm, baving Fig. 1. B. part of the Liver lying over it; its Figure somewhat resembling the usual make, inclining to that of an Half Moon; as appeared by that Hairy | Tophus we after. | Tab. 2. wards took out of it. But what was most remarkable Fig. 2. was, the Structure and Polition of the Two Orifices, (viz. that of the \* Gula, that leads into it, and the Pylo \* Tab. 2. rus t, that fends out) for they were both placed fo near Fig. 1. A. one another, that they feemed to touch or meet; and when I opened the Stomach, I found only a very flender Isthmus, or Wall, parted them. These Orifices were not at the Extreams of the Stomach, as usually; but inserted almost in the Middle of the upward Part, but more inclining towards that, that respects the | Duodenum. The | f. f. pouching or bagging + out at both Extreams, made it + D.D. somewhat resemble (as I said) an Half Moon. Stomach appeared but small, being much contracted, for it had not eaten any thing for some Days; it meafured about Three Inches and an Half in length, and about

about Two Inches in depth: The Gula which coveys the Food into the Stomach, confifted of strong Muscular Fibres, and was in all about Nine Inches in Length. The Pylorus, that carries out, seem'd to have its Passage free and open, without that annular Constriction or Valve, as in most other Animals; though here we observed a larger Body of Muscular Fibres, than in the other Intestines; which made me wonder how a Regurgitation of the Faces into the Stomach was prevented; but this we will consider anon.

But before we opened the Stomach, I observed at one Side a Perforation of Hole thorough, about the Bigness of an ordinary Pea, and Round. That it was occafioned by an Olcer there, I plainly perceived by the Lips or Edges; which were not fresh, but had an ulcerated Matter about them; and this, without doubt, was the Occasion of its Death; for it had fallen from its Food, and had pined away for some time before, and by its uneasy Motion, made its Keeper suspect, it had swallowed something that stuck in its Throat, or injured its Stomach.

A like Accident as this (as a Perforation of the Stomach)
I have Three Times met with in diffecting Human Bodies; and the last (being joined with another Physician in Consultation, a little before the Patient's Death) I foretold; and upon Diffection found confirmed. What appears to me, to be most likely to be the Cause of this Perforation, is, that some of the Glands in the Stomach (such as Payerus (qq) and Dr. Grew (rr) describes in the Intestines) being become Scrophulous or Steatomatous, might impossumate and so corrode the Coats of the Stomach, and cause a

D 2

Perfo-

<sup>(44)</sup> Exercetat. Anat. Med. de Glandulis Intestinorum. (17) Comparative Anatomy of Stomachs and Guts.

Perforation. And the rather I am of this Opinion, because in those Instances I mentioned of Human Bodies. I found in other Places of the Stomach, these Glands very large and Steatomatous; tho' Naturally they are but small, and often not observed. Where there is a Perforation of the Stomach upon an Inflammation, and upon that an Impostumation; there the Foramen is larger and not regular: as remarkably I once met with it in a Child. where a large part of one fide of the Stomach was Spha. celated. So likewife upon a Corrofive Poison taken, its Effects dilates it's felf more, and is not confined to fo narrow a Compass; as I observed once in one who had taken Rats-bane.

Upon observing this Perforation of the Stomach, I looked to fee, whether any of its Contents had been emptied into the Cavity of the Abdomen; but could find little or nothing: Nor indeed, when I came to open the Stomach, could I find any thing that could be evacuated that Way; for there was nothing contained in the Stomach, but a Body | of Clotted Hair, formed into the | Tab. 2. Shape and Figure of the Stomach, somewhat like an Half Fig. 4. Moon; covered with a flimy viscid Substance, which did serve the better to glue these Hairs together. These Hairy Tophi are frequently to be met with, in the Stomachs of Bruits, and I have had, and feen feveral, which have been taken out of Oxen; and the Butchers inform me, that they chiefly meet with them in the Winter Seafon, after the Hair begins to shed; and the Cattle feed upon Hay and dry Meat: But after the Spring, and in Summer, they do more seldom find them; as if the New Grafs, which Purges them, did contribute to diffolve these Tophi likewise. Georg. Hieron. Velscius has wrote Two Medico Philosophical Differtations (ff) about

<sup>(</sup>II) De Ægagropilis.

these Tophi, that are found in Goats; and others has made distinct Treatifes thereon; to whom I refer the Curious: and Gul. Pifo (tt) gives a Figure and Description of one.

But our Animal is Carnivorous, and in what all Accounts agree in, most Rapacious of the Winged Kind; and where it can't find its Prey on the Land, it will hunt for it in the Trees; most nimbly climbing them up: and if the tender Bough cannot bear the Weight of its Body on its Feet; by twisting its Tail about the Twig, it can hang thereby, and stretch it self the farther, to obtain its defired Food, or roba Nest. Nay, if I am not mis-informed, by this Means it can Fly, or pass from one Tree to another, without descending down; for thus hanging by its Tail, and waving and swinging its Body like a Pendulum, it can fling it felf into the Boughs of a Neighbouring Tree; where his Tail is fure to take fast hold of the first Bough it lights on, if otherwise it misses his Footing: and, as I have shewn, his hinder Legs being made like Hands, with a Thumb, it can more readily raise its Body up by them.

Noctambulum Animal est (laith de Laet (z) & cum cæteris Avibus, tum Gallinis insestissimum. Georg. Margrave (a) laith, Mordax est, vescitur libenter Gallinis, quas rapit, ut Vulpes, & Arbores scandendo, avibus institutur, vescitur quoq; sacchari cannis, quibus sustentavi per quatuor septimanas in cubiculo meo. So likewise Gul. Piso (b) Mordaces sunt ut Vulpes earumq; more Gallinis & Columnis non solum, sed Avibus in fastigiis Arborum institutur; quarum desestu Cannis sacchareis aliisq; optimis vegetabilibus vescuntur. So that they are not only Carnivorous, but when need drives them, they can take up

<sup>(11)</sup> Hift. Nat. & Med. lib. 5. p. 327.

with other Food. Scandit Arbores incredibili pernicitate (faith the Mexican History (c) diu in Cavernis latitat, vescitur Cobortalibus, quas Vulpecularum, Mustelarumq; sylvestrium more, jugulat, illarum sanguinem absorbens. And asterwards adds, Vivit enim in Calidis, ac pascitur Carne, fructibus, Pane, Oleribus, frumentaceis, aliisq; generibus, veluti nos experimento cognovimus, alentes iltud domi, ac in deliciis habentes. Which last Account seems true; for this that we dissected would eat any

thing, that was brought from the Table.

We shall now observe how the Food, when it is received into the Stomach, and contained there, till 'tis throughly digested, is afterwards dispensed with the greatest Advantage, for the Nourishment of this Creature. And what I hinted, how 'tis, that a Regurgitation of it into the Stomach again, is prevented; especially upon the Posture 'tis frequently in, when it hangs by its Tail, since (as I observed) the Passage at the Pylorus is so open and patent. And for the doing this, we must expect Nature's Contrivance (which is always admirable) to be great; not confining her self still to the same Rules; but is Insinite and All wise, in attaining the same Ends, with the greatest Variety and Mechanism.

Which leads me to consider the Structure and Order of the Intestines: But in doing this, I must first take Notice of the Mesenterie, that Membranous Part which colligates them, and fixes their Situation; and gives to them the Order of their Figure. For the Intestines are not just sastened to the Peripherie or outward Circumference of the Mesenterie; but the outward Membrane of the Mesenterie of both sides, is entirely projected and continued over the whole Canalis, or Duct, of the Guts; and is to them the outward or common Membrane: So that I have often, by separating this outward Membrane, from what lies under it, the Muscular; I have extracted

the whole length of the Guts; leaving only the Common Membrane, as tis continued from that of the Mesenterie; which I could inflate, as if the whole of the Guts remained.

Now here we observed that remarkable Difference, from what is in many other Animals; that we can't but make Two Mesenteries; one peculiar to the small Guts, the other belonging to the great ones, or Intestina Crassia, as they are called; for tho' continued to one another, yet the difference in their Figure or Bulk, is so much, that fully justifies the Distinction. And for Distinction sake, I shall call the former Mesenterium minorum, and the latter Mesenterium majorum, so. Intestinorum: for minora and majora I think more expressive, than tennia

and crassa; at least it appeared so in our present Subject.

If. f. For here I remarked, that as the || Duodenum descended from the Stomach, it ran under the + Colon, (just where 'tis joined to the Cæcum') towards the Middle of the Spine. Hence I found a Projection of the first Mesen-

Fig. I. u.

+ 1.66

w. terie || into a Spiral Line, like a Cochlea or winding Pair of Stairs: So that upon Inflation, these Intestines here, made several Convolutions, or Windings, the not exactly Spiral, but as represented in our Figure; and the better to shew the Currency of their Canalis here, and how those Gyrations follow each other; I have signified it by the Order of the Letters of the Alphabet; so that (g) is succeeded by (b) and (b) by (i) and so on: Not that I could represent the whole in this Figure; for some of these Gyri, at least great parts of them, lay dipt and hidden by others, that lay over them.

The Second Mesenterie +, or Mesenterium majorum, as I choose rather to call it, than Mesocolon (for it did belong likewise to the Cacum || and \* Reclum, as well as the Colon +) was projected more in a Plain; and made almost

most a Circular Figure at its Peripherie; so that the Cacum, and Colon did almost entirely encircle the fmall Guts.

The fmall Guts (as here inflated) measured about Six Feet and an half in length. The Caeum was about Six Inches long; and the Colon and Rectum Two Foot long. The Girth of the Duodenum (I mean all along here, as inflated) was Three Inches; the Ileon || Two In- || 4 ches and an Half; the Girth of the Caeum, in the largest Place, was Six Inches; of the Colon Four Inches; and the Rectum was Three Inches about. from the Spine to the utmost Projection of the fmall Guts, under the same Circumstance of Inflation, measured about Six Inches; the greatest Diameter that the Colon in this Circular Figure made, was somewhat above Seven Inches.

In the whole Dust or Canalu of the Intestines, I could not observe any Valves; no not at the Cacum its self. Tis true, that the Foramen into the Cacum, was a great deal less than the Capacity of the Gut its self; however, the Passage into it was so open and wide, as readily to receive or emit its Contents. For by pouring a large Quantity of Water into the Stomach, so as to wash out

would first run into the Cacum, if it was not filled before, and then into the Colon, and as readily, upon elevating the Colon, it would pass thence, first into the Cacum; and,

the Faces contained in the Intestines; I found, that it

when that was filled, into the Ileon.

Passing by other Remarks which I might make upon the Glands in the Inward Coat of the Intestines, upon the Blood-Vessels in the Mesenteries, &c. I shall now proceed to give my Thoughts upon the whole Structure and Figure of the Intestines, and how advantageously they are contrived for the Uses designed them by Nature.

For First, by means of these frequent Gyrations and windings of the Intestines, there is a greater Opportunity given to the Separation of the Chyle into the Vasa Chylistra; for the Bore of the Intestines being so large (as I have shewn) and there being no Valves in all their length; if there was not this Contrivance to give a Lett, or Remora to the hasty Descent of the Faces, they would pass off so nimbly, that with them a great part of the Chymous Substance likewise, would be carried off: But the Length of the Intestines here, being so great; and as they run, making so many Convolutions, they do prevent

this Danger.

Secondly, By means of this Cochlea, or Spiral Figure of the first Mesenterie, to which the small Guts are affixed, there may be prevented a Regurgitation of the Contents of the Intestines again into the Stomach, upon a Declivity of the Body of this Animal, as it is frequently in, when it hangs by its Tail. For tho', as I observed, the Passage from the Stomach, by the Pylorus, into the Duodenum, is large and open; yet in this Posture of the Body, there can't but be a Reduplication, or folding over of the Duodenum; fince the great Bulk or Wallet of these Intestines must incline and swag towards the Diaphragm; by which Reduplication, the Passage at the Pylorus must, in a great measure, be occluded; and the ascent of the Contents now, be altogether as difficult and great, as when the Animal stands upon its Four Feet.

The Reverse of this Structure of the Intestines I found and have described in my Anatomy (cc) of the Tajacu, or the Mexico Musk Hog: for here the Colon made a Spiral Figure (as I have there represented, in Tab. 1. Fig. 5.) and the small Guts made a Plain. In our Possum the small Guts makes a Spiral, and the Colon and great Guts a Plain. But a Spiral Convolution of the Intestines

is to be met with in several Animals, the their Structure be different; as in the Goat and Deer-Kind; and very

remarkably in a Woodcock.

But we will have done with the Guts, to proceed to other Parts; for their Comparative Anatomy is too large a Field, and would be too great a Digression, to engage in the Description of them in a single Subject.

The Pancreas was large, having one part (if I mis-remember not) running towards the Spleen, and the

other down by the Duodenum.

The Spleen was Two Inches and half long, and One Inch in the broadest Part, and was of a dark Red

Colour.

The Liver in this Animal was very large, of a bright Red Colour, confissing of Three Lobes †; Two of them † Tab. 2. were much larger than the third, which lay out of sight, \$\frac{Fig. 5.}{B.B.B.}\$ and was not to be seen, but upon inverting the Liver: and here we found not only at the Edges of one of the larger Lobes, deep Incisures \*, which rendered it jagged; \*eee but also in the middle of the Concave part of the same Lobe, several deep Fissures †: Possibly for this Reason, † d d d. that so it might yield and give way the better when 'tis inverted, as 'tis always, when this Animal hangs by its Tail. The Bladder \* of Gall here was very large. The \*C. Situation of the Liver and Spleen here, appeared as in other Animals.

In the Vrinary Parts I did not observe any thing peculiar or different from the usual Structure, unless what we shall remarke of the Bladder of Vrine. The Kidneys || || Tab. 2. of each side were a little above an Inch and half long, Fig. 2. AA. about three quarters of an Inch broad, and of the Figure almost of a Kidney-Bean. The Emulgent + Veins and Arte-+ bb. ries were very plainly seen: But on the inside of the Kid. \* cc. neys, towards the upper Part, were placed the Glandule Renales ||, or Renes Succenturiati, as they are called by || D. some,

fome, and which were here very large, and of the same Colour with the Kidneys themselves, which was a deep Red; whereas these Glandulæ Renales in Men and other Animals, are usually of a White, Yellowish Colour. The Treters \* were about five Inches and an half long, and were inserted into the Neck of the Bladder of Trine, as is represented !!, first running under, then ascending up by the two Extreams of each Tterw, as they lie duplicated. The Bladder + of Trine, being instated, was about the Bigness of a Hen's Egg and of that Figure. The Neck of the Bladder, or Trethra\*, (which was about an Inch long) lay over the Vaginæ Tteri +; and here the Trethra and the Vaginæ Tteri emptied themselves into one common Canalis \* or Passage, which measured about

an Inch and half in length.

I f.

+ G.

¥ b.

+ i i.

\* K.

In most Animals, about the Kidneys there uses to be observed a large Body of Fat covering them, being contained in the Membrana Adipofa: But here we found four large protuberant Lumps of Fat, two of each fide : two of them lying in the Pelvis of the Abdomen, near the Bladder of Urine, and the Vterine Parts; and the two others, between them and the Kidneys. Upon Examination, I found that they confifted of regular large Laminæ; which were eafily separable from one another, in broad Fleaks; fo as I have not observed before; and indeed, before I had examined these Lumps of Fat, I could not tell well what to make of them; and I am apr to think, that these Two Bodies of Fat, near the Vterus are, what Margrave (a) and Pifo (b) calls, Testes sub Ano interins jacentes; for, as I shall shew from themfelves, in the Male, the Testes are otherwise placed.

We shall proceed now to the Examination of the Vterine Parts: For 'tis so far from Truth, what is afferted by some, that it has no Vterus within, that here we find not only one, but two Vteri; and these too most

E 2

wonder-

wonderfully contrived, and far different from the common Structure and Make of this Part, in other Animals. And the more too I wonder at this Mistake, since they pretended to have diffected them; for lo Margrave (a). Hac Burfa (speaking of the Marsupium or Pouch) inse Vterus est Animalis; nam alium non babet, uti ex sectione illius Comperi ; in bac semen concipitur. & catuli formantur. And to Piso (b). Ex reiteratis borum Animalium Sectionibus, alium non invenimus Uterum præter banc Burfam ; in qua semen concipitur & Catuli formantur. But notwithstanding what they talk of their diffecting them. there is not one Observation (as I know) of any one of the inward Parts, that they have given us. Hernandez (c) is more in the Right (and indeed his Account is much more to be valued, and more faithful in the whole. than any of the others) where he tells us, Quaternos. quinofve parit Catulos, quos utero conceptos, editofq; in Lucem, Alvi capacitate quadam, dum adbuc parvuli sunt, claudit & lervat. We will therefore here take a Survey. and an Account of these Parts; and we find, that there are two Ovaria, two Tubæ Fallopianæ, two Cornua Uteri, two Uteri, and two Vaginæ Uteri.

The Ovaria || were placed one of each side, near the || f |.

Extreams of the Cornua Vieri, being fastened to the Ale \*\* rrr.

Vieri, and were about the Bigness of a Velch. The Va.

Sa Praparantia (the Arterie † and the Venie || that did go † n n n.

to and from them) were very plain, and as I have repre. || 0000.

sented them; though the greatest part of these Vessels

were bestowed upon the Cornua Vieri. Near the Ovaria,

I observed the Fimbria || Foliacea, and thence a Passage || Tab. 2.

into the Tuba Fallopiana \*. The Tuba Fallopiana \* were Fig. 3. b.b.

two fine slender Canales or Dusts, supported by the Ala Fig. 2. tr.

Vieri, and running waving, and led into the Extreams

of the Cornua Vieri. The Cornua † Vieri, being insta † Fig. 3.

ted, were about the Bigness of a Goose Quill, about an d. A. Fig. 2.

ted, were about the Bigness of a Goose Quill, about an d. M. n.

Inch and half long, and were fastened to the Ale Vieri, towards both Ends a little crooked, but where they pals into the Uteri, they were reflected inwards; at the other Extream reflected outwards. Their Substance seemed rather thicker than the Ziteri themselves, and not so transparent, by reason of the numerous Blood-Vessels which irrigated them almost all over; for in the infide, both above and under, there ran the whole length of the Gornua, large Trunks of Blood-Vessels, sending from the Sides all along numerous Branches; which is very requifite: For in Animals that are Multiparous, as is our Subject, the Litter or Fettu do lie, and are formed in the Cornua Uteri. And I did here take Notice of some little Risings of the inward Membrane of the Cornua, whereby they were somewhat divided into Cells; but very imperfectly: However, for the Nourishment and Formation of the Embrio's here, so great a Number of Blood-Vessels is highly necessary; and they were far more numerous here, than in the Vteri themselves. It has Four or Five Toung Ones at a time, faith Hernandez (c). Pife (b) and Joh. Stadius faith, Five or Six. That which Margrave (a) observed, had Six. Ralph Hamor (1) and Cardan lays, it has Seven. So Job. de Laer faith Six or Seven. But Jul. Cef. Scaliger (r) out-reckons them all; for he faith, Fecundissima est, duodenos parit exigues : But this Account I do suspect.

These Two Cornua do empty themselves into the Fig. 3. Two Uteri\*, just in the Middle, where they are considered together; and so outwardly seem to sorm, but as fig. 2. x x. it were, one continued Body †; from this Conjunction, near the Neck of the Bladder, extending themselves on each side, and afterwards, being resected to the Neck of the Bladder again, where they pass into the Vaginative it. But having extended this Part by Instation, and so letting it dry, and then disserting it; lobserved as

Mem

Membrane | like a Diaphragm, perfectly to run cross, and intirely to divide them, near the infertion of the Cornua, into two distinct Bodies; so that what is contained in the Vierus on the Right Side, can't pass into the Vierus on the Lest Side, by means of this Partition Wall; tho' outwardly (as I said) they both seemed, but as one conti-

nued Body.

I must consess, the Fabrick of this Part, seemed very surprising to me; and such as I have not met with a Parallel, or the like, in any Animal besides; at least of the Quadrupede Kind. 'Tis true, in Lobsters and Crabs, in the Female there are two Vieri, as in the Male there are two Penes, but more distinct and separated from one another. So two Penes, and each forked too, I have observed in the Rattle-Snake (un); but how the Male Possum is provided, I cannot tell: But this I think is the only Instance of a Land Quadrupede, that has two Vieri; and each of these too, seemingly double, by that Resection they make, and by an impersed Diaphragm, which divides the Cavity of each Vierus a considerable way, as we shall show.

These Vieri are not sastened to the Ale, as are the Ovaria, Tubæ and Cornua; but where they are conjoin'd near the Insertion of the Cornua, they do adhere very firmly to the Neck \* of the Bladder, not easily to be se. \* Fig.3.1 L. parated thence; and by Membranes to the Restum; where more separable. So that the Neck of the Bladder lies over that Diaphragm or Membrane which parted them (as I said) into two distinct Vieri. Here the Body of the Vieri seemed to be about the Bigness of the End of my Finger; or in Compass (thus instated) it measured about an Inch and three quarters: Hence they

<sup>(</sup>ww) Philosoph. Transact. No. 144.

were projected towards each fide, and not according to the Length of the Spine, gradually inlarging the inward Cavity, as 'tis extended. For here about the Angle of Reflection, it measured in Compass two Inches and an Half. The Uteri being thus extended towards each fide about the Space of an Inch and three quarters; and then reflected + back again, towards the Neck of the Bladder; and so pals into the two Vagina ||. which lies under the Fig. 2.11. Urethra \*. From this Angle of Reflection, the Cavity of each Uterus gradually lessens, and is much smaller than the other part of the Vierus. The Capacity of each Vierus being the largest at the outward Elbow, where it begins to be reflected; for here it made, as 'twere, one Common Cavity, for almost the length of an Inch: But on the infide, I observed a Membrane + to be projected from the internal fide of the Uteri, just from the Corner where the sides of the Uteri are doubled, whereby this Cavity is in part divided; and for this Reason, shall call this Membrane, the Second, or an imperfect Diaphragm of the Uteri.

t 7.

PPP.

Here in these Uteri, I observed Four large Trunks of Blood Veffels, which did run the whole length of them, lending from their fides numerous Branches, and Ramifications all along. These Trunks were propagated from the Hypogastrick | and Spermatic \* Vessels. I did also here observe in these Uteri (thus by Inflation extended and dried) several Fasciculi of Muscular Fibres, placed at a regular Distance from one another; which did run the whole Length of the Uteri likewise: by means of whose Contraction, the Fetus may be more easily forced out.

These two Uteri (as I mentioned) empty themselves tii. Fig. into the two Vagine t; for at this Extream, the U-3. f. teri, making a turn at the Neck of the Bladder, are continued thence into the two Vagina, which lie just under the

the Urethra\*, or that Passage or Pipe which conveys the \* Fig. 2. b. Urine from the Bladder, and are much of the same Length with that of the Urethra, which was about an Inch. Their Capacity was about the Bigness of a Wheat-Straw. Both thele Vagina and Urethra too, emptied themselves into a Common Passage ||, or Canalis, which I K. was as large as all the other Three; and about an Inch and half long: it looked Redish, by means of the numerous Blood-Veffels it enjoyed, and at last had its Exit so near the Fundament, that when alive, there was not observed any other Foramen outwardly, but that which led into the Redum. But when I came to diffect it, by elevating the Skin here, which seemed to cover it, like a Valve; I observed the Foramen that led into this Common Passage, and putting a Blow-pipe into it, at the same time, by Inflation I extended the Bladder of Urine, and the Uterine Parts too; viz. The Vogina, the Uteri, and the Cornua. So that in the Skin here, there was only one Foramen + + Tab. 1. for the Exit of the Faces, and the Urine and the Fatus Fig. 3. C. too.

I have had no Opportunity of diffecting a Male Poffum; and indeed, of none other but this fingle Subject: For had I, I might have been more exact in some Particulars; nor is it almost possible, to observe all in One.

The Account they give of the Male, is but very imperfect and short: Mas Faminae per omnia similis, bene testiculatus, saith Margrave (a) In Bursa pendulos Testes, more Gati, gerens. And much the same, saith Piso (b), Mas Famellae plane similis, in Bursa pendulos, more Gati, testiculos fert: But adds, Et quod notatu dignum, Manticam (licet à Famellae diversam) habet qua alternatis vicibus Catulos quoque circumfert. I could wish he had given us, wherein twas different; and what kind of Penis the Male had.

The Ingenious and most Learned Mr. Ray (o) Queries, Whether the Tai-ibi of Brafile, described by Margrave (a); differs from our Subject, the Possum, only in Sex? Or, Whether 'tis another Species of Animal? And indeed, by a Passage in his Description of it, one would think, that Margrave did take the Tai-ibi to be only the Male. I will transcribe the whole Paragraph, it not being very long, that every one may have the Liberty of making their own Conjectures. Tai-ibi Brafilienfibus, faith he) Lufitanis Chachorro do mato, Belgis een Boschratte: Animal corpore tereti & oblongo. Totius corporis cum collo longitudo ab occipitio ad caude initium quatuordecim digitorum, crassities decem. Caput habet vulpino æmulum, ore acuto, barba felina: oculos conspicuos & prominentes, nigros: aures subrotundas, molles, graciles, albas, teneras ut charta mollis. Crura, pedes & digitos cum unquibus habet ut fæmella jam descripta, uti & caudam. Totum corpus vestitum est pilis albis splendentibus, qui in extremitatibus nigricant, & magis quidem in dorso, maxime tamen in cruribus : circa anum & initium caudæ pene nigricant. Os & aures albicant. Cauda in exortu ad quinque digitorum longitudinem pilis vestitur albis in extremitate nigricantibus, reliqua pars major ad finem usque cincta est corio tenui squamoso albicante instar exuviarum serpentis. Fætet graviter, care tamen illius comeditur. Victitat libenter gallinis, ut vulpes. Pili inserti sunt tenui cuticulæ, que detrabi potest salvo manente corio crassiori. Testiculos propendentes habet ut felis mas. Had Margrave mentioned in his Description, the Marsupium or Pouch, it had been more clear; but by the Account he gives, one cannot but think, that he makes the Carigueya, the Female, and the Tai-ibi the Male of the same Species of Animal. As they eat the Tai-ibi, to Ralph Hamor (1) tells us, that he has eaten the Possum, and that 'tis a grateful and wholesom Food.

F

Pife (b) having concluded his Description of the Cariqueya, adds, In Indiis Orientalibus, idq; folum quantum hactenus constat in Amboina, similis Bestia frequens ad Falis magnitudinem accedens, mattata ab incolis comeditur. fi rite preparetur, nam alias fætet. Nomen illi Cous Cous inditum, What Similitude this Beaft may have with our Animal; or whether it is to be reckoned amongst the Animalia Crumenata, as Scaliger (r) calls them, I do not know. Our Animal feems to be properly a Native of America. Richardus Dinothus (as I find him quoted in Aldrovandus (a) faith Veram borum Patriam Americam effe, presertim Pervanis Regionibus. So Peter Martyr (w), Arbores in Pariana Regione ingentes funt, inter quas repertum est animal, &c. and then describes our Possum. That 'tis found in Darien and' Florida, is afferted by Nierembergius (d); and in New-Spain, by Cardan, and others. That they are in the Isles of Anguilla and Tabago, is affirmed by the Author of the Present State of his Majesty's Isles and Territories in America, pag. 138, and pag. 250. And in Virginia they are frequently to be mer with; as Ralph Hamor (1). John de Laet (m), Captain John Smith (n), and a great many others, tells us; and this that we diffected came from thence.

But not only in the West, but South America likewise, 'tis to be found: So Petrus Masseius (e) and Caspar Barlaus (f) assures us, that they are in Brasile. Whether Cardan (i) was not mis-informed, when he tells us, that 'tis in Æthiopia; I do very much question; his Words are these: Animal aliud mittit Æthiopia, parte anteriori vulpi persimile, Cauda & posteriore Cercopitheco, pedibus anterioribus bumanis, auribus vespertilionis, quod crumenam habet sub Ventre, qua Catulos undiquag; gerit nec dimittit, nisi dum lactare vult. What Authority he had for this, I do not know; but he owns too,

that they are in the West-Indies. Nierembergius (d) amongst the Places where this Animal is to be found. reckons likewise the Molucca Islands; which, if true, there may be something in what Pifo (b) saith of Amboina, which is one of them. But I think it does deserve a farther Enquiry.

I shall proceed now to give a Description of the See-

leton, and so shall conclude.

+ K.

1006

We will begin therefore with the Head, which, from the End of the Occiput, to the Exstream of the Nares, was Four Inches and Three Quarters long; of which the Rostrum + measured Three Inches; and just where Fig.4- AA the Rostrum and the Cranium || met, the Bones were so pinched in, at the Sides, that here, 'twas very narrow; and I may fay, in Proportion to the Bulk of the Animal, this was the least Cranium that ever I met with in a Quadrupede. On the Forebead, the Rostrum was an Inch broad, having on each fide, a Protuberance \* justing out. There was a large Sature + just in the Middle, which divided the upper Bones of the Nares lengthways, and though they ran flender towards the Extream of the Nares; yet these Bones towards the Forebead. spread into a Triangular Figure, and as they are joined together, they form a Rhomboide, or a Lozinge. But I will not be particular in describing each Bone, that compose the Head; for fear of being tedious: But I cannot but take notice of that remarkable rifing Ridge | like a Creft, that runs the length of the Cranium, from the Forehead to the Occiput, just in the Middle; where the Sutura Sagittalis is in other Skuls. This Ridge, for Distinction lake, I shall call, Protuberantia Offea longitudinalis; and I observed, it jutted out from the Cranium, above a Quarter of an Inch: Just at its upper Edge, I could perceive a Seam like a Suture; so that though now, these Bones

Bones are so well united together, that they appeared as one entire Body; yet in the Fætus, without doubt, they are separable, and are Two. And this I rather think, because in the upper part of the Cranium I could not find any Sutures at all. So likewise answerable to the Lamdoidal Suture, may be those other Ridges in the Extream of the Occiput, which I shall call, Protuberantia Offee Laterales \*; which arifing on each fide from the \* d. Processus Styloides, ascends obliquely up the hinder Part of the Occiput; and just in the Middle at the Top, is joined with the Longitudinal Ridge, I have described. These Ridges, although as deep as the first, yet were not standing so upright, but projected rather like a Pentbouse, over this hinder part of the Cranium; by both which Ridges, the Cranium is so well guarded and defended, that 'tis almost impossible, the Skull should be any ways cracked or broken. Something like these Ridges, but nothing so large, I have observed in the Skull of a Weafel.

And not only the Brain, but the Eyes likewise, are very well guarded and desended, by the Os Zygomaticum; which is very broad and strong; in the broad-+ e.f. eft Place being above Three Quarters of an Inch, and in the narrowest Half an Inch, being very thick on its under Edge; but at its upper, growing thin and sharp. But for the greater Strengthening this Bone (which is formed by a Process from the Os Temporum ||, and ano-|| e. ther from the Maxilla superior \*) where they meet, they + f. lap over one another, and so become the stronger. This Os Zygomaticum was Two Inches and an Half long, and standing off from the Cranium an Inch in Distance.

In the Orbit of the Eye at the inward Cantbus, there was a large Foramen +, which led into the Cavity of the Nose, and by a Dust placed here, the Tears or Moisture from the Eyes is conveyed into the Nostrils. In the up-

b. per Jaw Bone likewise, there was a large Foramen ||, which was for the Passage of some Vessels from the inward Orbit of the Eve.

The Court

The Cranium +, which encompais'd the Brain, in the largest Place, was about an Inch over; and about an Inch and Half in Length; but its Cavity jutted out somewhat farther towards the Nares, making as it were, a particular Cell here, and pretty capacious, for the receiving the Processus Mamillares, and that fore Part of the Brain. And afterwards I observed the Os Cribriforme very remarkably perforated with Holes, like a Sive; and indeed, in forming this Organ of Smelling, Nature feems very careful and follicitous, the Rostrum making so great a part of the Head, that the Cranium it felf feemed very inconsiderable in respect to it, its inward Capacity containing not above the Quantity of a Walnut. Not having diffected the Brain (which I hope I may have hereafter an Opportunity to do) I will not at present, insist on the Description of the several Foramina's I observed on the inside, for the Passage of the Nerves and Blood Veffels. But, as I have thewn, fince it has fo little Brains, Nature hath been very kind to it, in preferving them, by the Defence of those Prominent Bones I have remarked, both length-ways and laterally behind; and on the Sides too, 'tis guarded by the Temporal Process + of the Os Zygomaticum.

The Os Spongiosum in each Nostril, seemed very curiously contrived, by the abundance of Lamina it enjoys; so that the Membrane that covers them, by this means, is render'd more Capacious, and capable of receiving more plentifully the Effluvia's of those Animals, it would either catch, or avoid; and in this Sensory'tis known, that Bruits excel even Man himself, and their

Organ is more adapted for it.

The Under Jaw | confished of Two strong Bones, join- 1 1. ed together only at the Mentum; each measured Four Inches in Length. The Head of this Bone (which was Half an Inch broad) was received into a Simus of the Os Temporum, and very firmly articulated there. It had Two Processus: the Anterior † or Superior is large and + m. thin, into which is inserted the Temporal Muscle. The Inserior \* Process is smaller, and runs to a sharp Point: \* n. Here at this Process, the Edge of the Mandible is so dilated, that it measured above Half an Inch. On the instide of the Jaw here, is a large Sinus, which leads to a Foramen that goes into the Body of the Jaw-Bone, and

affords a Passage for the Vessels thither.

The Use of these Bones is for Mastication; which leads me to confider the Teeth. And here we find all the Three Sorts of them; for in the Upper-Jaw before, were Eight small Dentes Incifores, Four of each Side; then a void Space, almost a Quarter of an Inch; then Two large Prominent Dentes Canini, one of each Side; which jutted out of the Faw about Half an Inch : these were Succeeded of each Side, with Three Dentes Incifores; but these were much stronger and larger than the Fore-Teeth; and these imitated the Dentes Molares, in that they were inserted into the Jaw-Bone with Two Phangs: But the Heads of these Incifores were acuminated; whereas the Heads of the Molares were flat, and almost of a Triangular Figure. There were Four Dentes Molares of each Side: in all, Four and Twenty Teeth in the Upper-Faw. But the double Phangs of the Molares, and the Incifores Majores were such, as at first fight, one would think them Two distinct Teeth; each Phang being inferted into a distinct Alveolus, or Socket in the Jaw, and remaining separated some way above the Jaw Bone, and only joined at the Head.

In the Under Jaw-Bone, there were likewise of each Side, Four Dentes Incisores Minores before; then a little void Space; after that, the Dens Caninus; then Three Dentes Incisores Majores; and last of all Four Dentes Molares, answerable to those in the Upper Jaw, but somewhat smaller. In both Jaws in all, Eight and Forty Teeth. Georg. Margravius's (a) Account I find is somewhat different from mine; for he saith, Dentes (babet) instar falis seu Vulpis, anterius nimirum tam superius, quam inserius parvulos, binc quatuor longos Caninos, superiores quidem longiores, inseriores breviores, binc iterum sex alios, & dein Molares. Nimirum sedecim Molares, duodecim inter medios, quatuor Caninos, & incisores parvulos in inseriori mandibula octo, in superiori decem, nameduos majusculos in medio habet, more Leporis.

But we will proceed next to what the Head is fastened to, the Vertebra: and I find here, Seven Vertebra of the Neck; Thirteen of the Back or Thorax; Six of the Loins; Three of the Os Sacrum; and Two and Twenty of the Tail; One and Fifty in all, and all extraordinari-

ly contrived.

The First Vertebra t of the Neck (to which the Head? † No. 1. is fastened, and is therefore called the Atlas) had Two broad transverse Processes, but no Spine. The Second Vertebra \* of the Neck, had a very large and thick Tab. 2. Spine of a Triangular Figure; and in it was observed a Fig. 6. b. large Semi-circular Sinus , which was so deep, as to re-¥ 6. ceive into its Bosom, a great part of the First Vertebra; by which means, the Articulation was very much strengthened. This Vertebra is called Dentata, from that Tooth like Protuberance + I have represented, and which + 4. is received into the Hollow of the First Vertebra, where the Medulla Spinalis runs. This Vertebra backwards, had Two Processus obliqui superiores , and Two Oliqui. inferiores t. The Third Vertebra of the Neck, had the + f.

fame.

fame Processes || both before and behind; but the Spine \* | Fig. 7. here was about Three Quarters of an Inch in Height; " diff. about the Third of an Inch thick; and just at the Top feemed to be a little cleft. The Fourth and the Fifth Vertebra had the same Processes, as the Third Vertebra; and the Spine here, likewise very thick, and cleft at the Top; but gradually lessening in Height, as also Thickness. The Sixth Vertebra, besides the former Processes, had likewise an acute Transverse one, on each Side; and its Spine much shorter, and more acuminated than the for-The Seventh Vertebra of the Neck had only Two oblique Processes before, and none behind; and Two acute transverse Processes, and a very short and sharp Spine: So that upon holding up the Head, the Spine of the First Vertebra + of the Thorax, would touch the + Tab. 1. Top of the Fifth Vertebra of the Neck. These Verte- Fig.4. No.2. bre are fo strongly and closely locked into one another, that though each of them are large in themselves; yet, thus arriculated, they do not make full Two Inches in But What I most wondered at, was, to ob-Length. ferve the Thickness and Strength of the Spines of the Second, Third, Fourth and Fifth Vertebræ, far different from what I have observed in any other Animal; and can't but think must be, for the better defending its self from the Injuries it might receive by any Fall; and to perform much the same Office, as that Prominent Bony Ridge in the Cranium. And not only here, but likewife the Spines of several of the Vertebræ of the Thorax and Loins, I find are thus flatted, and broad at the Ends; and they being fo, should it happen to fall to the Ground, by Chance or Defign; these Spines being so Prominent and flat, will better bare off the Blow; and upon this Account, there is no Danger of his breaking his Neck, his Back, or his Head; fince they are all Three so well fecured and guarded.

The first Seven Vertebra of the Thorax, have Two + 74b. 2. oblique Processes & forwards , which run under the Fig. 8. 46 hinder oblique Processes of the preceding Vertebra ; and have Two oblique Processes \* backwards, which rides \* c c. over those of the succeeding Vertebra; as likewise Two transverse Processes to which at their Ends have small + d d. Acetabula's | or Sinus's, for the receiving the Heads of le c. the Ribs, which are fastened to them. The Spines of ¥ 4. these Vertebra, are flender, thin, and sharp; about Three Quarters of an Inch long, The Six following Vertebra of the Thorax, have thort, thick, and flat Spines. The oblique Processes being continued on each Side of the Sping, make as 'twere, a Gutter; and the transverse Processes here, are somewhat different from the former, The Spines | of the Vertebra of the Back 1 Tab. 2. or Loins, the more they approached the Os Sacram, for Fig. 9. they lessened gradually in their Thickness on the Edge. + bb. cc. But here were double oblique Processes +, viz. Four at each End of the Vertebra, and the undermost spreading \* Tab. 1. themselves out broad. The Three Vertebre \* of the Os Fig. 4. Sacrum, are firmly fastened to the Os Ilium +; but the No. 10. + No. 13. last not so entirely as the Two former: But this at each Side had a broad transverse Process, and the Spines of \* No. 11. thefe were thin. The Two First Vertebra t of the Fail had only One (mall acute Spine; but in all the other Wertebra of the Tail, both at the Head and Tail of each Vertebra, I observed Two Spines; but those at the Head of the Joint, the larger. In the Six First Vertebra of the Tail, there was, of each Side, a broad transverse Proceft, the Length of the Joint: In the other Vertebre only at the Head and Joil, a jutting out at the Sides. The Vertebre about the Middle of the Tail, were the longeffe being there about an Inch long; nearer the Root of the Tail, and at the End not fo long. Brengthering the whole, there was observed your Dialeles

G

But

But what I was most of all pleased to see, and I think. is a wonderful Piece of Nature's Mechanifm, was, those Spines + or Hooks placed in a Line, in the Middle of the + No. 12. under Side of the Vertebra of the Tail. Tis true, the 12. first Three Vertebre had none of these Spines, nor were they necessary here, fince they lay within the Compass of the Offa Coxendicis; but in all the other Merrebre, to the End of the Tail, they were to be observed But as they approached the Extream of the Tail, they grew leffer and thorter. Thele Spines | (where longest) were about | Tab. 2. a Quarter of an Inch, or somewhat more: they were Fig. 10. bbb. placed just at the Articulation of each Joint, and as P faid, in the Middle from the Sides and keemed to be articulated, both to the preceding and following Pertebra; not being an entire folid Body, but saifing from the Vertebra with Two Legs or Crura, become afterwards perfectly united at the Ends. By this means, thefe Bones are rendred more firm and ftrong, and this Hollow . . . . ferves for the transmitting the Blood-Veffels thorough them; and one may observe here a Stria, or Furrow, all the Length of the Vertebra, for the receiving them; whereby they are the better fecured from Compression, when this Animal hangs by his Tail. And for the performing this Office, nothing, I think, could be more advantageously contrived: For when the Tail is twirld or wound about a Stick, this Hook of the Spine eafily fustains the Weight, and there is but little Labour of the Mulcles required, only enough for the bowing or crooking the Tail; for then, as by a Hook, the Weight of the whole Body is hereby suspended. And for the doing this, 'twis' observed, that in each preceding Vertebra, there did a Muscle arise, which was inserted on each fide of the fucceeding Vertebra; which Acting or Contracting, must necessarily bend and curve that Joine. But for the strengthening the whole, there was observed Four Muscles

to arise from the Os Sacrum, which did run the whole length of the Tail; Two on the upper Side; and Two on the under; fending each a Tendon to each Internale or Kercebran So that when the Skin was ftript off, the outward parts of these Muscles seemed to have tendinous Expansions over them; the whole Length of the Tail. and almost to be covered by them; which must needs very much contribute, and add Strength to the Tail: belicks what may be the Effect of their Infertion of Tendons into each Joint, or Vertebra, in curling and un-

bearing the Tail in this all woll among character when alive, and how advantagiously 'tis contrived for that Purpole. we have sufficiently feen; But I find it is highly commended by Hernandez (c), as an extraordinary Medicine and Specifich; and from him, several others relate the lame : I thall give you only his Words, and to proceed on to our Sceleton; Canda bujus Animalu (faith he) egregium elt Medicamentum; trita enim drachma unius menlura, atq; ex agua aliquoties devorata, nullo præassumpto alimento co die quo bauritur, Urine meatur mire abstergit. eadem epocata, traffifg lapillis, & quacunque alia re meatus obstruente. Venerem excitat, generat lac, ac perfractis medetur, & Chalicis, Partum accelerat, Menfes elicit, extrabita; tufa & imposita aculeas qui sunt infixi, ac ventrem emollit. And concludes, Et fortaffis mullum extat Medicamentum ad bas res amnes peragendas, pre-

! Tab. 2. Fig. 4.

To the Vertebra of the Thorax are fallened the Ribs !. and there are Thirteen of each Side. The Seven foremost are more persocily articulated with the Sternum; the Six fucceeding may be reckoned in some Senfe, Coffe Nothe: For though they are long, and as they proceed from the Versehre, are inclined backwards, towards the hinder Legs; yet afterwards they are reflected for-

forwards towards the Sternum of Cartilago Scuttformis. But I must here take Notice, that though in Man, and other Animals, that Part of the Ribs that is fastened to the Os Pectoris, or Sternum, be usually Careilaginous; yet here, in our Subject, I observed it to be all Bony throughout. However, this Difference I found, that the Ribs did look redder, by reason of the Blood Vessels in them; and this Part was Whiter, and where it was fastened to the Ribs, one might plainly see; so that it may well pals for a Bony Cartilage; as often, the Cartilages do become Bony. The First Rib was only an Inch long, and its Bony Cartilage a Quarter of an Inch: hence gradully the Ribs increase in Length; for the Seventh Rib was Three Inches long; and its Cartilage One Inch and Half. The Four last of the Coffe Nothe, gradually lessen again in Length; for the last Rib of all was only One Inch and Three Quarters long; and its Cartilage did not run Home to the Os Pettoris, or Sternum, though the First, Second and Third of the Colla Notha did. orics devorata, nuita praca

The Os Pedoris, or Sternum, confisted of Seven Bones, according to the Number of the Fore-Ribs, that are fastened to them. At the Beginning of the Sternum, there jutted out a sharp Bony Cartilage, which, from its Figure, I shall call, Cartilage Enfiformis +; and here + p. was fastened One Extream of the Clavicule ||; at the End of the Sternum, towards the Belly, there was a broad, roundish Cartilage, which therefore I shall call, Cartilage Scutiformis \*.

There were Two Clavicule || or Collar-Bones, each an Inch and Half long; having one Extream fastened to the First Bone of the Sternum, or the Cartilago Enfformis; and the other End to the Spine † of the Scapula, near † r. the Conjunction of it to the Os Humeri. By means of this Bone, it can more advantagiously bring its Fore-Feet

(653)

to its Month; as it uses to do when it feeds its felf, as do the Monkey-Kind, who have Clavicula too as well as Man; though many Animals want these Bones.

The Scapula or Shoulder-Blade was about Two Inches long, about an Inch and Half broad; its Spine though thin, yet the nearer it approached the Shoulder, it grew larger and flatters. Into the Simus of the Neck of the Tab. 1. Scapula, was received the Head + of the Shoulder-Bone, Fig. 6.4.

or of the Fore-Thigh-Bone; as to that Protuberance, called, the Acromium, was fastened the End of the Clavicula.

This Thigh-Bone of the Fore-Legs I found very re| Tab. 1. markable; so that I caused a Figure | to be made of it:
| Fig. 6. 'twas about Two Inches and Three Quarters long; 'twas
| thick and Strong, having a large rough Spine ' jutting
| forward, and running Halt the Length of it. The low| tf. 8. er Extream † of this Thigh-Bone, to which was fastened

the Tibia and Fibula, grew very broad, being almost an Inch broad. Above, where this Bone began to grow broad, on the out-fide, was a large Protuberance |; and on the Inside there was a great oblong Foramen\*, or hol-

low Passage, formed by a small Bone arising from the inward Fore Part of the Thigh Bone, where it begins to grow larger, and was afterwards united to that part of the Basis + of this Bone, where the Fibula, or Minus

focile is joined. Just in the Middle of the Basis of this Bone, there was a large Sinus || which backwards appeared deeper, which did lock into another deep Sinus of the Tibia; by which means these Bones were so firmly articulated together, as they were not easily, if possibly, to be put out of Joine.

The Tibia †, or Focile majus, was a strong Bone, about Three Inches long; which was extended upwards about a Quarter of an Inch above its Articulation with the Thigh Bone; and at the other End, was fastened to the outward Bone of the Tarsus.

The Fibala, or Focile Minus, was a smaller Bone, w. w. placed more inward and forwarder, and nor to long as the Tibia; being articulated above (but not to firmly) with the Thigh Bone, and below, with the inward Bone of the Tarfus. For there were but Two Bones of the Tarfus, having each a small Simu, for the receiving taxable Heads of the Two Fociles. The Bones of the Metalenafus were Four, or it may be Five to which were 19 joined the Five Fingers or Toos of the Fore Feer. The innermost Toe had but Two Articulations or Joints, but at the End had a large booked strong Nail: The other Four Fingers had each, Three Articular or Joints, armed with Hooked Nails, as the First and the land of the Mails.

The Hinder-Legs were lastened to the Trunk of the Body by the Os Innominatum; which, though properly here is but Two Bones, (viz. One of each Side) being separated above, by the Os Sacrum, and below, are joined together at the Or Pubis ) yet commonly they do, for Distinction take; give Three Names to each; as the uppermost Part of this Bone they call, the Or Ilium + + 1 No. 43. the lowermost, the Or Hebii, or Coxendicis |; and the Fig. 5. 66. fore Part, where the Coahtion is, the Os Pubis " The + a a. Length of the whole, in a ftraight Line, was Three Inches. In the Os Ifchii was the Acetabulum to being at ff. large Socket, for the receiving the Head of the hinder Thigh Bone; and deeper in, there was a Space for the fastening the Ligament; from which Space, there was a Sinus which led outward; fo that the Brims of the Acetabulum was not an entire Circle, but broken off here. But the most remarkable Bones here, are, the Offa Mar-Supialia \*, seu fanitores Marsupii, of which I have given + co. a Description before; therefore now shall proceed.

The Hinder-Thigh Bone was a little above Three Inches long; 'twas roundith, and a strong Bone. But the Tibia +, + Tab. v. or Majus Focile of the Hinder Leg, was somewhat Fig. 4.TT.

longer, and a little curved. The Fibula |, or Minus Focile, was about the same Length, Braighter and Senderer a This, towards the Foot, was articulated to the Os Calcis; as the Tibia was to to the Talus, or Aftragalus; and these Two Bones I make the Tarfus ; and joining + > >. to them, were the Bones of the Metatarfus +; and to thefe 1 7. the Phalanges | of the Fingers or Toes. In the innermoft, or the Thumb, there were only Two Articuli, or Rones:

in the other Four Toes, or Digiti, in each there were Three Articuli, or Joints. The End of the Thumb was more flatted, than the Ends of the other Toes: For the Thumb, as I have observed, bad a flat Nail, like a Human Thumb; in the others, the Nails were long, and curved. I observed likewise, at the Articulation of each Foint of the Toes, on the under Side, there were Two Imall Bones, that are called, Offa Sefamoidea, and thefe, both in the Fore and Hinder Feet.

Banks of fome River

lam fensible how tedious I have been, in the Defeription of this one Animal, and from a fingle Observation too! Had I had more Leasure to have drawn it up in. I might have been much shorter. But it being an Animal, fo very remerkable; and one too, fui Generis, or a distinct Species from all others, I was the more inclin'd to be as particular as I could, though not fo much as I could have wished. And I am the more confirmed in what I have formerly wished (ww) that for the perfecting a Natural History of Animals, we had a distinct Account and Anatomy of some One of a Species; which, with a little Variation, might ferve for all of that Family: fince I find so great a Master of Natural History, as Mr. Ray (xx), " as I have foyed them at play we noming omal and lo it

<sup>(</sup>www) In my Preliminary Discourse concerning Anatomy, and a Natural History of Animals, before my Phocena. (ax) Synophy Animal p. 324.

Since part of this Discourse was Printed off, I have met with a Paffage in a Treatife of the Bucaniers of Ameriea, wrote first in Dutch by John Equemeling, and fince translated into English (yy), which I can't but take Notice of: And if true, (as the Author affures us, that he 49. has feen it often) 'cis an Instance of a Quadrupede, which I knew not of before, that receives its Toung into its Belly. But'ris not a Land, but Water Animal; or it may be rather, one Amphibious, between both. 'Tis a fort of Crocodile, which he calls a Cayman; his Words are thefe, which I therefore repeat, that by farther Observation. if true; it might be confirmed; as also, it might be more particularly specified, whether their Entrance into the Belly, was by the Mouth, or any other Part. " Ma-" ny times (faith he) their Eggs (for 'tis an Animal Ovi-"parous) are destroyed by Birds, that find them out, as " they scrape amongst the Sands: Hereupon the Females of the Caymans, at fuch times as they fear the coming " of any Flocks of Birds, do oft-times, by Night, Iwallow "these their Eggs, and keep them in their Stomach till " the Danger is over. And from time to time, they bu-" ry them again in the Sand, as I have told you, bring-" ing them forth again out of their Belly, till the Season " is come, of being excluded the Shell. At this time, " if the Mother be nigh at hand, they run unto her, " and play with her, as little Whelps would do with their "Dams, sporting themselves according to their own " Cuftom. In this fort of Sport, they will often times " run in and out of their Mothers Belly, even as Rabits " into their Holes. This I have feen them do many times, " as I have spyed them at play with their Dam, over the "Water, upon the contrary Banks of some River: at " which time I have often dilturbed their Sport, by throwat ing a Stone that way, causing them on a sudden to creep " into the Mother's Bowels, for fear of fome eminent a Danger. The

## EXPLANATION of the FIGURES.

FIGURA PRIMA

here vained to the Olia

R Epresents the outward Shape and Figure of the Possum, drawn from the Life,

FIGURA SECUNDA
Represents the Slit or Aperture in
the Belly that goes to the Marsuprum or Pouch, where the
Toung Ones lodge, till they can
shift for themselves.

FIGURA TERTIA.

A. Shews the Marsupium or Pouch turned the inside ontwards where may be observed the Hair or Fur that covers it, and may help the better to keep the Toung Ones warm.

BB. The two binder Legs cut off.

C. The Foramen of the Anus, which is also the common outward Vent or Exit to the Rectum, the Bladder of Urine, and the Uteri too.

D. The beginning of the Tail.

Gives a View of the Sceletanior Bones of this Animal.

23. The Rollrum or Smout.

bb. The Cranium or Skull that did contain the Brain.

ccc. A Bony Ridge or Protuberan-

8. The Stormach.

tia Offea longitudinalis, that did run the length of the Cranium, and over a part of the Roftrum. d. The lateral Ridge, which, like a Penthonfe, justed over the bin?

of the I this in the Fore

der pare of rhe Cranium, Pro-

e.f. The Os Zygomaticum (e) its Process from the Os Temporum, and (t) that from the Maxilla Superior, or Upper Jawa

g. A Foramen or Hole in the inward Canthus of the Orbit of the Eye that leads into the Noffrils, and by a Duck conveys the Tears or Moisture of the Eyes into Them.

h. A Foramen or Hole in the upper Jam for a Passage to the Vessels. i.A Protuberance of the Os Frontis.

It. A Sature of the Os Nariuro . 1

m. The Superior Process of the un-

The inferior Prices of the under

o. The Clavicula of one Side.

p. The Carrilago Enfitormis of the first Bone of the Strengen,

q. The Scapula or Shoulder-Blade Bone. A sile guira M all O

H side 1 10 to The

r. The Spine of the Scapula.

SSSS. The Thigh-Bones of all the Feet.

TTTT. The Tibia, or Focile majus of all the Feet.

uu. Part of the Tibia in the Fore-Legs, extended beyond the Articulation.

www. The Fibula or Focile minus in all the Legs.

xxxx. The Bones of the Tarfus,

yy.yy. The Bones of the Metatarlus. zzzz. The Toes.

as The Thumbs in the binder Feet. No.1. The first Vertebra of the Neck called the Atlas

2,3,4,5,6,7. The fecond third fourth. fifth fixth, and feventh Vertebra of the Neck.

8. The first Vertebra of the Thorax. 9. The first Vertebra of the Loins.

10. The first Vertebra of the Os so Sacrum. slot to m

Coxygis, or Tail.

12.82.12.42. The Spines or Hooks the infide of the Tail.

13,14. The Os Innominatum, where (13.) is the Os Ilium, (14) the Os Ischii or Coxendicis.

15.15. The Offa Marsupialia, Sen Janitores Marfupii.

\* \* \* \* The Ribs, thirteen in all.

(i). The Cartilago Scutiformis. FIGURA QUINTA Represents the Situation of the Ossa Marsupialia, &c.

The Offa Pubis.

b. The Coalition or the joining of the Offa Pubis.

cc. The two Offa Marsupialia, or Ja-

nitores Marsupii.

d,e. The Bafis of the Offa Marfupialia, where joined to the Offa Pubis, (d) the inward Head of the Basis, (e) the outward.

f.f. The Acetabulum or Socket for receiving the Head of the Thigh Bone.

g.g. The Os Ileum.

hi. The Vertebræ of the Os facrum. II. The Os IIchii or Coxendicis.

FIGURA SEXTA

Exhibits the Figure of the fore fide of the Thigh Bone of the fore

a. The Head of the Thigh Bone, where tis fastened to the Scapula.

b. A large rough Spine, which runs above balf the Length of this Thigh Bone.

11. The first Vertebra of the Os C. A Protuberance of this Bone on the out fide.

> d. A large Foramen or bollow Paffage.

> e. A Sinus for receiving the Head of the Tibia.

> f.g. The Bafts or lower Extream of the Thigh Bone.

TABULA SECUNDA. FIGURA PRIMA

Represents the Stomach and Guts, and the several Coylings they make.

A. The Gula or Gullet.

B. The Stomach.

canfed by an Olcer there.

dd. The two pouching out of the Stomach at the two Ends.

e. The Pylorus.

f. The beginning of the Duodenum. g,h,i,k,l,m,n,o,p,q. Represents the Small Guts, and the Coyles and Convolutions they do make. Some of the Coyles lie bid, and out of fight; But the order bom the follow one another, is figuified by the Order of the Letters of the Alphabet : fo that (g) follows (f), and (g) is succeeded by (i) and (i) by (k), and fo on to (q) where the Ilion is discharged and emptied into the Cæcum or, if that is full, into the Co-

RR. The Cæcum. SSS. The Colon.

T. The Rectum.

V. The first Mesenteric, or Mesenterium minorum Intestinorum

lon at the first Letter S.

W. The fecond Melenterie, or Melenterium majorum Intestinorum.

FIGURA SECUNDA Exhibits the Urinary and the U terine Parts.

AA. The two Kidneys.

bb. The Emulgent Veins.

cc. The Emulgent Arteries.

dd. The Glandulæ Renales.

ce. The two Ureters.

f. The Infertion of the left Ureter, into the Neck of the Bladder.

c. A Perforation of the Stomach, G. The Bladder of Urine turn'd afid

h. The Urethre of a lati well of

ii. The two Vaginz Uteri.

K. The common Paffage from the Urethra, and the two Yaginte. L The Arteria Aorta, or Great Ar-

terie.

m. The Vena Cava.

nn.n. The Spermatick Arteries. 00.00. The Spermatick Veins.

p.p.p. The Hypogastick Arteries and Veins.

err. The Ala Uteri, fen potins Cornuum.

ss, The Ovaria.

tt. The Tube Fallopiane.

uu. The Cornu Uteri of the Left Side opened.

w. The Cornu Uteri of the Right-Side not apened.

xx. The two Uteri opened.

y. The Diapheagm that divides the two Utori.

zz. The imperfett Diaphragma, which partly divides each Uterus, and lies over the Paffage of that Part of the Uterus, which is doubled and tends to the Vaguz: 132 A FI DI

FIGURA TERTIA More particularly shows the Uterine Parts.

AA. The two Ovaria.

bb. The Fimbria Foliacea.

oc. The Tubæ Fallopianæ,

dd. The two Cornua Uteri.

EE. The two Uteri reduplicated

1 A Slie in the Neck of the Left Orerus | d Spinals pefferio woith volve & A to flow its Paffage into the Vagina on that Side.

g. The Left Vagina opened.

h. The Oftium or Mouth of the Right Vagina.

1. The common Paffage from the Urethra and Vaging. .51Y91

m. The Vega Cavandra V adT .m

II. The Bladder of Urine cut off.

EIGURA QUARTA Represents the Hairy Tophus, or Ball of Hair that was taken out of the Stomach.

FIGURA QUINTA Represents the Liver. Congum.

A. The Vena Cava. The O sale 22

BBB. The three Lobes of the Liver,

C. The Bladder of Gall. ddd. The Fissures in the Body of the Liver cce. The Incilures at the Edges of the Liver,

FIGURA SEXTA. A. The Spine of the fecond Verrebra of

the Neckago real coursed ... 2X b. Represents its Thickness.

c. A large Sinus for the receiving the first Verrebra.

d. The Dens or Tooth of this Vertebra. e. The Processus obliques superior of one Side.

f. The Processus obliques inferior of aa. Two Vertebras of the Tail; - L the fame Side bus boldnob zi

FIGURA SEPTIMA.

A. Represents the Spine of the third Vertebra of the Neck, where is

(hewn its natural thickness. b. The Hole through which the Medulla

cc. Two Small Boremina for the Pof-Sage of Veliels.

d. Represents the Cleft at the top of the Spine.

ee. The two Processus obliqui Supe-The beginning of therefoldering

Fif. The rue Processus obliqui. Inferiores before ; has

Represents the first Verrebra of the Thorax.

A. The Spine, which is long and seute.

bb. The Oblique Processes before. cc. The Oblique Proceffes behind.

dd. The Transverse Processes.

ce. Where the Ribs are fastened

f. The Hollow where the Medulla Spinalis paffes.

FIGURA NONA Represents the fourth Vertebra of the Loins.

aa. The two upper Oblique Processes bebind.

b. The Spine.

cc. The two under Oblique Processes bebind.

FIGURA DECIMA Represents the second and third Vertebra of the Tail hugas ad

bbh The Spines or Hooks on the infide, by means of which, it can better bang by its Tail.

cc. A Hollow or Foramen in the middle of these Spines, through which,

Blood Veffels paffes.

## ERRATA.

PAS. 1. 1. 5. r. at Chirurgions p. 2. 1. 21. r. Simi-vulpa. p. 3. 1. 26. r. Epithet. p. 4. f. 1. r. determined. 1. 14. r. addidi. p. 5. 1. 7. r. Crumenam. p. 6. 1. 8. r. her. p. 3. 1. 6. r. Hexagons. p. 9. 1. 21. r. funk. p. 10. 1. 32. r. invikem. p. 12. 1. 5. r. proficialtur. 1. 10. r. quaritare didicerint. 1. 13. r. funk. p. 10. 1. 22. r. invikem. p. 21. 1. 12. r. fra. p. 29. 1. 29. Columbia. p. 36. 1. 25. r. Vetch. 1. 26. r. Vein. p. 42. 1. 2. r. felis. p. 43. 1. 10. r. Extream. 1. 19. r. Sature. p. 44. 1. 5. r. Lambdoidal. p. 45. 1. 1. r. Scive. p. 47. 1. 9. r. felis.